arXiv:1212.3399v2 [math.NT] 8 Feb 2013

THE AUTOMORPHY OF CERTAIN $K3$ FIBERED CALABI–YAU THREEFOLDS WITH INVOLUTION OVER $\mathbb{Q}$

YASUHIRO GOTO, RON LIVNÉ, AND NORIKO YUI

Abstract. We consider certain $K3$-fibered Calabi–Yau threefolds over $\mathbb{Q}$ with involution. We will discuss the automorphy of the Galois representations associated to these Calabi–Yau threefolds. We construct such Calabi–Yau threefolds as the quotients of products of $K3$ surfaces S and elliptic curves. We choose $K3$ surfaces S over $\mathbb{Q}$ with non-symplectic involution σ acting by -1 on $H^{2,0}(S)$. We fish out $K3$ surfaces with involution σ from the famous 95 families of $K3$ surfaces from the list of Reid [18], and of Yonemura [21] where Yonemura described hypersurfaces defining these $K3$ surfaces in weighted projective 3-spaces.

Our first result is that for all but few (in fact, nine) of the 95 families of $K3$ surfaces S over $\mathbb{Q}$ in Reid–Yanemura list, there are subfamilies of defining hypersurface equations which are of Delsarte or Fermat type and endowed with non-symplectic involution σ. One implication of this result is that with this choice of defining equation (S,σ) becomes of CM type.

Let E be an elliptic curve over $\mathbb{Q}$ with involution ι, and let X be the standard (crepant) resolution, defined over $\mathbb{Q}$, of the quotient threefold $E\times S/\iota\times\sigma$, where (S,σ) is one of the above $K3$ surfaces over $\mathbb{Q}$ of CM type. One of our main results is the automorphy of the L-function of X.

The moduli spaces of these Calabi–Yau threefolds are Shimura varieties. Our result shows the existence of a CM point in the moduli space.

Contents

1. Introduction 2
2. $K3$ surfaces with involution 3
3. Construction of the Calabi–Yau threefolds 5
4. Realization of $K3$ surfaces as hypersurfaces over $\mathbb{Q}$ 6
5. Computations of Nikulin's invariants for $K3$ surfaces of Borcea type 10
6. Computations of Nikulin's invariants for $K3$ surfaces of non-Borcea type 17
7. Realization of Calabi–Yau threefolds as hypersurfaces over $\mathbb{Q}$ 21
8. Singularities and resolutions on Calabi–Yau threefolds 23
9. The automorphy of Calabi–Yau threefolds over $\mathbb{Q}$ 27
10. Base change and automorphic induction 29
11. Proof of the automorphy of our Calabi–Yau threefolds over $\mathbb{Q}$ 31
12. The automorphy of mirror Calabi–Yau threefolds over $\mathbb{Q}$ 33
13. Tables 36
References 37

Date: February 11, 2013.

1991 *Mathematics Subject Classification.* Primary 14J32, 14J20; Secondary .

Key words and phrases. Calabi–Yau threefolds, involutions, $K3$-fibrations, Elliptic fibrations, CM points, Shimura varieties, Calabi–Yau varieties of CM type, L-series, automorphy, motives.

1. Introduction

We will address the automorphy of the Galois representations associated to certain $K3$-fibered Calabi-Yau threefolds over $\mathbb{Q}$. Here by the automorphy, we mean that the L-series of the Calabi-Yau threefolds over $\mathbb{Q}$ come from automorphic representation in $GL_n(\mathbb{A}_{\mathbb{Q}})$ for some n. In fact these represetations arise by automorphic induction from GL_2 using a generalization of the work of Arthur–Clozel [1].

Our Calabi–Yau threefolds were previously considered by Voisin [20] and by Borcea [4] from the point of view of geometry and also towards physics (mirror symmetry) applications.

We now describe briefly the Voisin–Borcea construction of Calabi–Yau threefolds over $\mathbb{C}$. Let E be any elliptic curve with an involution $\iota : x \mapsto -x$, and let S be a $K3$ surface with involution σ acting by -1 on $H^{2,0}(S)$. The Hodge structure of the product $E \times S$ is given by the tensor product of the Hodge structures of the components. The quotient threefold $E\times S/\iota\times\sigma$ is singular, but the singularities are all cyclic quotient singularities, and there is an explicit crepant resolution, which yields a smooth Calabi–Yau threefold X. The Hodge numbers $h^{1,1}(X)$ and $h^{2,1}(X)$ are determined by the fixed locus S^σ of σ on S.

To find our $K3$ surfaces, we use the famous 95 families of $K3$ surfaces which can be given by a weighted homogeneous equation in weighted projective 3-space, as classified by M. Reid [18] (see also Fletcher [7]). Yonemura [21] gave explicit equations for these surfaces as weighted hypersurfaces $h(x_0, x_1, x_2, x_3) = 0$ using toric methods, and we will use his list throughout this article.

We first fish out from the list of Yonemura subfamilies S of $K3$ surfaces having a convenient involution σ acting on the holomorphic 2-forms of the surfaces as multiplication by -1. Borcea [4] found 48 such pairs (S, σ) in Yonemura's list. We find additional $41 + (3)$ such pairs (S, σ) in Yonemura's list (our involutions have a different formula from Borcea's examples), bringing the total to 92 pairs.

Nikulin [16] classified all $K3$ surfaces (S, σ) over $\mathbb{C}$ with non-symplectic involution σ by triplets of integers (r, a, δ), and found that there are 75 triplets up to deformation. In this paper, we calculate only the invariants r and a for our 92 examples and realize at least 40 triplets (r, a, δ) of Nikulin. Since $\delta \in \{0, 1\}$, this number may increase somewhat. However, as the task of calculating δ is more involved, especially because we often need a $\mathbb{Z}$-basis for $\mathrm{Pic}\,(S)$, we leave the determination of the invariant δ to a future publication(s).

For 86 of our 92 pairs of (S, σ) we find a representative hypersurface defining equation for S of Delsarte type over $\mathbb{Q}$, that is, the equation consists exactly of four monomials with rational coefficients (the other terms are chosen as 0). Toric methods show that the chosen equation defines a $K3$ surface if it satisfies the following conditions: (i) for any i, $0 \leq i \leq 3$, it must contain terms of the form x_i^n or $x_i^n x_j$ $(i \neq j)$ with non-zero coefficients; (ii) it must be quasi-smooth (cf. Section 4) and (iii) it has the same singularity configuration as the original hypersurface. (We should mention why we only have 86 pairs: This is because for the six weights, $K3$ surfaces have involution but cannot be realized as quasi-smooth hypersurfaces in four monomials.)

Once this condition is satisfied, we obtain $K3$ surfaces S of Delsarte, hence of CM type. By definition, a cohomology group of a variety is of CM type if its Hodge group is commutative; a variety is of CM type if all its cohomology groups are of CM type (see Zarhin, [22]). In general the computation of Hodge groups is notoriously difficult. However, Zarhin's results (see Livné-Schütt–Yui [13, Theorem 8] for a convenient statement) give an indirect way for our cases, using the argument developed in [13]: a Delsarte surface S can be realized as a quotient of a Fermat surface by some finite group, and one knows since Weil that Fermat (hyper)surfaces are of CM type.

It is known [4], [19] that over $\mathbb{C}$ the moduli spaces of Nikulin's $K3$ families are Shimura varieties. Recently, the rationality of the moduli spaces of all but two out of the 75 Nikulin's families of $K3$ surfaces has been established by Ma [14, 15], combined with the results of Kondo [11], and Dolgachev–Kondo [6].

Our results give explicit CM points in these moduli spaces defined over $\mathbb{Q}$; we do not know what their fields of definition (or moduli) are in the Shimura variety.

Next we take a product $E \times S$, where S is a $K3$ surface of CM type over $\mathbb{Q}$ with involution σ as before, and E is an elliptic curve over $\mathbb{Q}$ with the -1-involution ι. Let X be the Calabi–Yau threefold obtained as the crepant resolution of the quotient of the product threefold $E \times S/\iota \times \sigma$. Then we show that X has a model over $\mathbb{Q}$. We can establish the automorphy of the Galois representations associated to X. We show that all the cohomology groups of X are of CM type if and only if E also has complex multiplication. This parallels the work by Livné and Yui [12] on the modularity of the rank 4 motive associated to the Calabi–Yau threefolds over $\mathbb{Q}$ obtained from the quotients of the products $E \times S/\iota \times \sigma$ where S are singular $K3$ surfaces with involution σ.

We also construct mirror partners $X^\vee$ (if they exist) of our Calabi–Yau threefolds using Borcea–Voisin construction. In fact, 57 of the 95 families of $K3$ surfaces S of Reid and Yonemura have mirror partners $S^\vee$ within the list. We show that all these 57 families have subfamilies with involution σ and a CM point rational over $\mathbb{Q}$. Then the quotients of the products $E \times S^\vee/\iota \times \sigma$ give rise to mirror partners of $E \times S/\iota \times \sigma$.

From the point of view of mirror symmetry computations, our results supply particularly convenient base points in both the moduli space and in the mirror moduli space: they are defined over $\mathbb{Q}$, and their cohomology Galois representations are attached to automorphic forms whose L-functions are known.

2. $K3$ SURFACES WITH INVOLUTION

Let S be a $K3$ surface over $\mathbb{C}$. Then $H^2(S,\mathbb{Z})$ is torsion-free and the intersection pairing gives it the structure of a lattice, even and unimodular, of rank 22 and signature $(3,19)$. Therefore, by the classification theorem of such lattices, up to isometry,

$$H^2(S,\mathbb{Z}) \simeq U^3 \oplus (-E_8)^2$$

where U is the usual hyperbolic lattice of rank 2 and E_8 is the unique even unimodular lattice of rank 8.

Let $\mathrm{Pic}\,(S)$ be the Picard lattice of S. It is torsion free and finitely generated, and together with the intersection pairing it can be identified as the sublattice $\mathrm{Pic}\,(S) = H^2(S,\mathbb{Z}) \cap H^{1,1}(S)$ of $H^2(S,\mathbb{Z})$. We define the transcendental lattice of S, denoted by $T(S)$, to be the orthogonal complement of $\mathrm{Pic}\,(S)$ in $H^2(S,\mathbb{Z})$, i.e., $T(S) := \mathrm{Pic}\,(S)^\perp$ in $H^2(S,\mathbb{Z})$, with respect to the intersection pairing.

Consider now a pair (S,σ), where S is a $K3$ surface and σ is an involution of S acting by -1 on $H^{2,0}(S)$. Let $\mathrm{Pic}\,(S)^\sigma$ denote the part of $\mathrm{Pic}\,(S)$ fixed by σ. Let $(\mathrm{Pic}\,(S)^\sigma)^* := \mathrm{Hom}(\mathrm{Pic}\,(\mathrm{S})^\sigma,\mathbb{Z})$ be the dual lattice of $\mathrm{Pic}\,(S)^\sigma$. Let $T(S)_0 =$



Figure 1. Nikulin's pyramid

$(\mathrm{Pic}\,(S)^\sigma)^\perp$ be the orthogonal complement of $\mathrm{Pic}\,(S)^\sigma$ in $H^2(S,\mathbb{Z})$, and let $T(S)_0^*$ be the dual lattice of $T(S)_0$. From the assumption that σ acts as -1 on the holomorphic 2-forms of S one can show that it acts by -1 on $T(S)_0$ (and by 1 on $\mathrm{Pic}\,(S)^\sigma$).

Consider the quotient groups $(\mathrm{Pic}\,(S)^\sigma)^*/\mathrm{Pic}\,(S)^\sigma$ and $T(S)_0^*/T(S)_0$. Since $H^2(S,\mathbb{Z})$ is unimodular, the two quotient abelian groups are canonically isomorphic:

$$(\mathrm{Pic}\,(S)^\sigma)^*/\mathrm{Pic}\,(S)^\sigma \simeq T(S)_0^*/T(S)_0.$$

On $(\mathrm{Pic}\,(S)^\sigma)^*/\mathrm{Pic}\,(S)^\sigma$, σ acts by 1, while on $T(S)_0^*/T(S)_0$ it acts by -1. Then only for the group $(\mathbb{Z}/2\mathbb{Z})^a$, -1 is equal to 1. This shows that

$$(\mathrm{Pic}\,(S)^\sigma)^*/\mathrm{Pic}\,(S)^\sigma \simeq (\mathbb{Z}/2\mathbb{Z})^a \quad \text{for some non-negative integer } a.$$

Nikulin [16, 17] has classified such pairs (S,σ).

Theorem 2.1 (Nikulin). *The pair (S,σ) of $K3$ surfaces with non-symplectic involution σ is determined, up to deformation, by a triplet (r,a,δ), where $r = \mathrm{rank}\,\mathrm{Pic}\,(S)^\sigma$, $(\mathrm{Pic}\,(S)^\sigma)^*/\mathrm{Pic}\,(S)^\sigma \simeq (\mathbb{Z}/2\mathbb{Z})^a$, and $\delta = 0$ if $(x^*)^2 \in \mathbb{Z}$ for any $x^* \in (\mathrm{Pic}\,(S)^\sigma)^*$, and 1 otherwise.*

There are in total 75 triplets (r,a,δ), as shown in Figure 1.

The moduli space of (S,σ) with given triplet (r,a,δ) is a bounded symmetric domain of type IV having dimension $20-r$.

For a given pair (S,σ) of a $K3$ surface S with involution σ, we now consider the geometric structure of the fixed part S^σ of S under σ (i.e., the part where σ acts as identity). We follow Voisin [20] for this exposition.

Proposition 2.2. *There are three types for S^σ:*

(I) For $(r, a, \delta) \neq (10, 10, 0)$, $(10, 8, 0)$, S^σ is a disjoint union of a smooth curve C_g of genus g and k rational curves L_i:

$$S^\sigma = C_g \cup L_1 \cup \cdots \cup L_k$$

(II) For $(r, a, \delta) = (10, 10, 0)$, $S^\sigma = \emptyset$.

(III) For $(r, a, \delta) = (10, 8, 0)$, S^σ is a disjoint union of two elliptic curves C_1 and $\bar{C}_1$:

$$S^\sigma = C_1 \cup \bar{C}_1.$$

Furthermore, in the case (I), the genus g and the number k of rational curves can be determined in terms of the triplet (r, a, δ) as follows:

$$g = \frac{1}{2}(22 - r - a),$$

and

$$k = \frac{1}{2}(r - a).$$

Remark 2.1. Since σ is a non-symplectic involution, the quotient S/σ is either a rational surface or Enriques surface. It is an Enriques surface if and only if $S^\sigma = \emptyset$, i.e. $(r, a, \delta) = (10, 10, 0)$. On the other hand, if σ is a symplectic involution, then the minimal resolution of S/σ is again a $K3$ surface.

3. Construction of the Calabi–Yau threefolds

In this section, we recall the Voisin–Borcea construction of Calabi–Yau threefolds.

Let E be an elliptic curve with the standard involution ι, and let (S, σ) be a pair of $K3$ surface S with involution σ acting by -1 on $H^{2,0}(S)$. By the classification theorem of Nikulin, the isomorphism class of such pairs (S, σ) is determined by a triplet (r, a, δ) as we discussed in Section 2.

Now we consider the product $E \times S$, and the quotient threefold

$$E \times S/\iota \times \sigma.$$

Obviously, this quotient is a singular, having cyclic quotient singularities. We resolve singularities to obtain a smooth crepant resolution, denoted by $X = X(r, a, \delta)$, which is a Calabi–Yau threefold.

Proposition 3.1. [Borcea [3]] *The Hodge numbers of the Calabi–Yau threefold $X = X(r, a, \delta)$ are determined by the given triplet (r, a, δ), or equivalently, by the data from S^σ: Indeed,*

$$h^{1,1}(X) = 5 + 3r - 2a = 1 + r + 4(k + 1),$$
$$h^{2,1}(X) = 65 - 3r - 2a = 1 + (20 - r) + 4g$$

where k, g are described in Proposition 2.2. The Euler characteristic of X is

$$e(X) = 2(h^{1,1}(X) - h^{2,1}(X)) = 2(r - 10).$$

[Voisin [20]] *Put $N := 1+k$ and $N' := g$. That is, N is the number of components, and N' is the sum of genera of components, of S^σ. Then*

$$h^{1,1}(X) = 11 + 5N - N',$$
$$h^{2,1}(X) = 11 + 5N' - N$$

and the Euler characteristic is

$$e(X) = 2(h^{1,1}(X) - h^{2,1}(X)) = 12(N - N').$$

Now we discuss briefly resolution of singularities; detailed discussions are in Section 6. As above, let $\iota : E \to E$ be the standard involution. The fixed part E^ι consists of four points $\{P_1, P_2, P_3, P_4\}$.

We consider the generic case (I) when the fixed part S^σ of S is given by

$$S^\sigma = C_g \cup L_1 \cup L_2 \cup \cdots \cup L_k$$

where C_g is a smooth curve of genus $g \geq 1$ and L_i is a rational curve for each $i = 1, \cdots, k$.

Proposition 3.2. *The quotient threefold* $E \times S/\iota \times \sigma$ *has singularities along* $\{P_i\} \times S^\sigma$ $(i = 1, 2, 3, 4)$. *Each singular locus is a cyclic quotient singularity by a group action of order* 2.

By resolving singularities, we obtain a smooth Calabi–Yau threefold X*:*

$$E \times S/\iota \times \sigma \leftarrow X.$$

The exceptional divisors are four copies of a union of ruled surfaces

$$S^\sigma \times \mathbb{P}^1 := (C_g \times \mathbb{P}^1) \cup (L_1 \times \mathbb{P}^1) \cup \cdots \cup (L_k \times \mathbb{P}^1).$$

4. REALIZATION OF $K3$ SURFACES AS HYPERSURFACES OVER $\mathbb{Q}$

We are interested in finding defining equations over $\mathbb{Q}$ for pairs (S, σ) of $K3$ surfaces S with non-symplectic involution σ. For this, we appeal to the famous 95 families of $K3$ surfaces of M. Reid [18] (see also Fletcher [7]) and of Yonemura [21]. All these 95 families of $K3$ surfaces are realized in weighted projective 3-spaces $\mathbb{P}^3(w_0, w_1, w_2, w_3)$. Reid determined 95 possible weights (w_0, w_1, w_2, w_3), and singularities as they are all determined by weights. Then Yonemura described hypersurfaces defining them, using toric constructions.

We first recall the results of Borcea [4] and of Yonemura [21]. Here we say that $Q = (w_0, w_1, w_2, w_3)$ is *normalized* if $\gcd(w_i, w_j, w_k) = 1$ for every distinct i, j, k. Also, we assume that w_i's are ordered in such a way that $w_0 \geq w_1 \geq w_2 \geq w_3$.

Proposition 4.1. (A) [Borcea] *Assume that* $Q = (w_0, w_1, w_2, w_3)$ *is normalized and* $w_0 = w_1 + w_2 + w_3$. *Then there are in total* 48 *weights* (w_0, w_1, w_2, w_3) *giving rise to pairs* (S, σ) *of* $K3$ *surfaces* S *with involution* σ *acting by* -1 *on* $H^{2,0}(S)$. *More precisely, if* w_0 *is odd, there are* 29 *weights, and if* w_0 *is even, there are* 19 *weights.*

(B) [Yonemura] *Under the same assumption as in (A), a pair* (S, σ) *may be realized as the minimal resolution of a hypersurface* S_0 *of degree* $2w_0$ *in* $\mathbb{P}^3(w_0, w_1, w_2, w_3)$ *of the form*

$$x_0^2 = f(x_1, x_2, x_3)$$

where $\deg(x_i) = w_i$ *for* $0 \leq i \leq 3$, *the involution* σ *on* S_0 *is defined by* $\sigma(x_0) = -x_0$, *and* f *is a homogeneous polynomial in the variables* x_1, x_2, x_3 *of degree* $2w_0$.

Our first result is to extend the list of Borcea and Yonemura by adding more weights that yield $K3$ surfaces (S, σ) with involution σ.

Theorem 4.2. *There are in total* $92 = 48 + 44$ *weights* (w_0, w_1, w_2, w_3) *giving rise to pairs* (S, σ) *of* $K3$ *surfaces* S *with non-symplectic involution* σ *defined by* $\sigma(x_i) = -x_i$ *for some single variable* x_i*. In other words, we have* 44 *new weights (i.e., not in the list of Borcea) yielding* $K3$ *surfaces with involution* σ*.*

We divide the 92 *cases into two groups:*

(i) *The* 48 *weights of Borcea and defining equations for quasi-smooth* $K3$ *surfaces* S_0 *are tabulated in* Tables 1, 2 *and* 3 *in* Section 13.

(ii) *The additional* 44 *weights and defining equations* $F(x_0, x_1, x_2, x_3) = 0$ *for quasi-smooth* $K3$ *surfaces* S_0 *are tabulated in* Tables 4, 5, 6 *and* 7 *in* Section 13.

Proof. The proof of Theorem 4.2 is done by case by case analysis. Yonemura [21] determined hypersurfaces defining $K3$ surfaces using toric geometry. We use his list of equations to find $K3$ surfaces with non-symplectic involutions.

If the defining equation contains the term x_0^2 or $x_0^2 x_i$, then we can define involution σ by $\sigma(x_0) = -x_0$ just as in Borcea's 48 cases. If the defining equation contains the term $x_0^2 x_i + x_0 x_j^m$ (say), then we remove $x_0 x_j^m$ to define the involution $\sigma(x_0) = -x_0$ (see Tables 4 and 7).

For the equations in Table 5, we change x_0^3 to $x_0^2 x_1$ to define an involution by $\sigma(x_0) = -x_0$.

For the equations in Table 6, we choose variables other than x_0 (and remove several terms if necessary) to define an involution.

Note that in each of the 92 cases, the quotient S/σ is a rational or Enriques surface. Hence σ is a non-symplectic involution. $\square$

Remark 4.1. Among the 95 $K3$ weights of Reid, there are three cases #15, #53, #54 where we find no obvious involution; that is, there is no involution σ on S acting as $\sigma(x_i) = -x_i$ for some variable x_i. These cases are tabulated in Table 8 in Section 13.

For our arithmetic purposes, we like to have equations of *Delsarte type* (i.e. equations consisting of the same number of monomials as the variables) named after Delsarte (see [13], Section 4). Hypersurfaces of Delsarte type are finite quotients of Fermat varieties. Our next task is to find the subfamily of the 92 cases of Theorem 4.2 that satisfy the following conditions:

1. By removing and/or changing some terms of the original defining equation $h(x_0, x_1, x_2, x_3)$ of S in [21], we obtain a new equation of Delsarte type. In other words, the new surface S is defined by an equation consisting exactly of four monomials.
2. The new hypersurface obtained in (1) is quasi-smooth.

Conditions (1) and (2) give a restriction on the form of S. They change the arithmetic properties of S, but many of its geometric properties are unchanged. For instance, the types of singularities on S remain the same as the original hypersurfaces $h(x_0, x_1, x_2, x_3) = 0$ of [21].

Theorem 4.3. *There are* 86 *weights* (w_0, w_1, w_2, w_3) *that give rise to pairs* (S, σ) *of* $K3$ *surfaces* S *of Delsarte type with non-symplectic involution* σ*.* S *is the minimal resolution of a quasi-smooth* $K3$ *surface* S_0 *defined by a Delsarte equation in* $\mathbb{P}^3(w_0, w_1, w_2, w_3)$*.*

(a) *If* (S,σ) *is one of the* 48 *pairs determined in* Proposition 4.1 *other than* $\#90, \#91, \#93$, *then* S_0 *can be defined by an equation over* $\mathbb{Q}$ *of four monomials*

$$x_0^2 = f(x_1,x_2,x_3) \quad \subset \mathbb{P}^3(w_0,w_1,w_2,w_3).$$

The equation is obtained by removing several terms from the equation of Yonemura [21], *where* f *is a homogeneous polynomial over* $\mathbb{Q}$ *of degree* $w_0+w_1+w_2+w_3$ *(cf.* Tables 1, 2, 3*).*

(b) *Let* (S,σ) *be one of the additional* 41 *pairs determined in* Theorem 4.2 (ii) *other than* $\#85, \#94, \#95$. *Then* S_0 *can be defined by an equation* $F(x_0,x_1,x_2,x_3) = 0$ *over* $\mathbb{Q}$ *consisting of four monomials of degree* $w_0+w_1+w_2+w_3$. *In most cases,* $F(x_0,x_1,x_2,x_3)$ *can be chosen as*

$$F(x_0,x_1,x_2,x_3) = x_0^2x_i + f(x_1,x_2,x_3).$$

The weights and equations are listed in Tables 4, 5 *and* 6 *in* Section 13*.*

Proof. This can be proven by case by case checking of the list of equations of Yonemura [21]. In both (a) and (b), we transform S into a Delsarte type by removing several terms of its original defining equation. In doing so, we make sure that the condition (2) above is satisfied so that the new surface is also quasi-smooth.

Condition (2) is met if for each variable x_i $(0 \le i \le 3)$, the set of monomials containing x_i takes one of the following forms:

$$x_i^n,\ x_i^nx_j,\ x_i^n + x_ix_j^m,\ x_i^nx_k + x_ix_j^m$$

for some j and k $(j \neq k)$ different from i.

This choice for the defining hypersurface preserves the configuration of singularities (i.e., types and the number of singularities) on S as the original hypersurfaces. The point is that for each of the 86 families, we can always find a subfamily whose defining equation consists of four monomials containing terms x_i^n or $x_i^nx_j$ $(i \neq j)$ with nonzero coefficients. □

Remark 4.2. Our list of defining equations for S do not cover all possible equations of Delsarte type. For instance, in case #19 of weight $(3,2,2,1)$ in Table 4, we may also consider an equation $x_0^2x_1 + x_1^3x_2 + x_2^4 + x_3^8 = 0$.

Remark 4.3. For the 95 families of quasi-smooth weighted $K3$ hypersurfaces, Yonemura [21] described the number of parameters (i.e., the number of monomials) for their defining equations. The minimum number was four, but often equations contain more than four monomials. Our result shows that except for the six cases $\#85, \#94, \#95$ of Table 7 and $\#90, \#91, \#93$ of Table 1, the minimum number of parameters is attained.

Example 4.4. • Consider #42 in Yonemura =#3 in Borcea. Weight is $(5,3,1,1)$ and a hypersurface is given by

$$x_0^2 = f(x_1,x_2,x_3) = x_1^3x_3 + x_1^3x_3 + x_2^{10} + x_3^{10}$$

of degree 10. We can remove the second monomial $x_1^3x_3$. The singularity is of type A_2.

• Consider #78 in Yonemura = #10 in Borcea. Weight is $(11,6,4,1)$ and a hypersurface is given by

$$x_0^2 = f(x_1,x_2,x_3) = x_1^3x_2 + x_1^3x_3^4 + x_1x_2^4 + x_2^5x_3^2 + x_3^{22}$$

of degree 22. We can remove the second and the fourth monomials $x_1^3x_3^4$ and $x_2^6x_3^2$. The singularity is of type $A_1 + A_3 + A_5$.

- Consider #19 in Yonemura. Weight is $(3, 2, 2, 1)$ and a hypersurface is given by

$$F(x_0, x_1, x_2, x_3) = x_0^2x_1 + x_0^2x_2 + x_0^2x_3^2 + x_1^4 + x_2^4 + x_3^8$$

of degree 8. We can remove the first or the second monomials $x_0^2x_1$ or $x_0^2x_2$, the third monomial $x_0^2x_3^2$. The involution is given by $x_0 \to -x_0$. The singularity is of type $4A_1 + A_2$.

Remark 4.4. The cases #85, #90, #91, #93, #94 and #95 of Yonemura cannot be realized as quasi-smooth hypersurfaces in four monomials with involution σ.

For instance, consider the case #85 of weight $(5, 4, 3, 2)$ and degree 14. All the possible monomials of degree 14 are

$$x_0^2x_1 + x_0^2x_3^2 + x_0x_1x_2x_3 + x_1x_2^3 + x_0x_2x_3^3 + x_1^3x_3 + x_1^2x_2^2$$
$$+x_1^2x_3^3 + x_1x_2^2x_3^2 + x_1x_3^5 + x_2^4x_3 + x_2^2x_3^4 + x_3^7.$$

To make it quasi-smooth and defined by four monomials, we remove the terms $x_0^2x_3^2$, $x_0x_1x_2x_3$, $x_0x_2x_3^3$, $x_1^2x_2^2$, $x_1^2x_3^3$, $x_1x_2^2x_3^2$ and $x_2^2x_3^4$. The polynomial becomes

$$x_0^2x_1 + x_1x_2^3 + x_1^3x_3 + x_1x_3^5 + x_2^4x_3 + x_3^7.$$

Now no matter what two monomials we remove from it, it cannot be made quasi-smooth.

Note that if we allow more than four monomials, we can define a non-symplectic involution on this surface. For example, the surface defined by

$$x_0^2x_1 + x_0^2x_3^2 + x_1^3x_3 + x_1^2x_3^2 + x_2x_3^5 + x_2^4x_3 + x_3^7 = 0$$

is quasi-smooth and endowed with an involution $\sigma(x_0) = -x_0$.

Theorem 4.5. *Let (S, σ) be one of the pairs of* Theorem 4.3. *Then (S, σ) is defined over $\mathbb{Q}$, and there exists a quaduple (ρ, K, ι, χ) with the following properties:*

1. *ρ is an (Artin) Galois representation of Gal$(\overline{\mathbb{Q}}/\mathbb{Q})$,*
2. *K is a CM abelian extension of $\mathbb{Q}$,*
3. *$\iota : K \to \mathbb{C}$ is an embedding,*
4. *χ is a Hecke character of K of ∞ type $z \to \iota(z)^2$,*
5. $\dim \rho + [K : \mathbb{Q}] = 22$

such that the L-function of $S/\mathbb{Q}$ is given by

$$L(S/\mathbb{Q}, s) = \zeta(s)\zeta(s + 2)L(\rho, s + 1)L(\chi, s)$$

Proof. We follow a similar argument in [13], to which we refer for further details. Since S is a surface defined over $\mathbb{Q}$, its $\mathbb{Q}_\ell$-cohomology is 1-dimensional in dimensions 1 and 4, which contribute the factors $\zeta(s)$ and $\zeta(s + 2)$ respectively. Since S is a $K3$ surface the first and the third cohomology groups vanish, giving no contribution to the L-function. The second cohomology group is a direct sum $N_S \oplus T_S$ of the algebraic part,spanned by the subspace N_S of algebraic cycles and its orthogonal complement T_S, called the space of transcendental cycles. This direct sum decomposition is Galois invariant. Since N_S is the image by the cycle map of the Néron-Severi group NS (S) (with scalars extended to $\mathbb{Q}_\ell$), the Galois group acts on NS (S) through a finite quotient. Hence it acts on N_S by the Tate twist $\rho(1)$ of the corresponding Artin representation ρ.

To obtain the last factor we first consider the cohomology with complex coefficients. Since the defining equation for S uses only 4 monomials, it is a Delsarte surface. Hence it is a quotient of a surface in $\mathbb{P}^3$ with a (homogeneous) diagonal equation $\sum_{i=1,\dots,4} a_i w_i^r$ by some diagonal action of roots of unity. Moreover in our case the monomials in the (diagonal) equation for S have coefficients 1, which implies that the $a_i's$ can also be taken to be all 1's. Weil's calculation (see [13], Section 6) gives that over an appropriate cyclotomic field the Galois representation on the transcendental cycles is a sum of 1-dimensional representations coming from Jacobi sums, of infinity type as in the statement of the Theorem, which the absolute Galois groups permutes transitively. The Theorem follows. (For detailed discussion on Jacobi sums, see Section 10 below.) $\square$

5. Computations of Nikulin's invariants for $K3$ surfaces of Borcea type

Recall that a $K3$ surface S with involution σ is determined up to deformation by a triplet (r, a, δ), where r is the rank of $\mathrm{Pic}\,(S)^\sigma$. In this section, we compute r and a for $K3$ surfaces of Borcea type. By Proposition 2.2, r and a can be computed through the fixed locus S^σ.

Lemma 5.1. *Let $S^\sigma = C_g \cup L_1 \cup \cdots \cup L_k$ be the disjoint irreducible decomposition of S^σ, where C_g is a smooth curve of genus g and $L_1, \cdots, L_k$ are rational curves. Then*

$$r = 11 - g + k,$$
$$a = 11 - g - k.$$

Direct computation of r often requires a basis for $\mathrm{Pic}\,(S)$ or at least for $\mathrm{Pic}\,(S) \otimes \mathbb{Q}$. Since $\mathrm{Pic}\,(S)$ is usually difficult to determine, the fixed locus S^σ is often easier to handle than the Picard group.

In what follows, first we explain a general algorithm of computing g and k (and hence r and a). To describe the algorithm in detail, we choose $K3$ surfaces of Borcea type, namely those defined by $x_0^2 = f(x_1, x_2, x_3)$. After that, we explain how to compute r directly by looking at the σ-action on $\mathrm{Pic}\,(S)$ (see Theorem 5.5). It has a merit that we can obtain a closed formula for r.

Algorithm for the computation of g and k

We explain how to compute g and k (and then r) for our $K3$ surfaces S. In this section and the next, we choose S_0 to be a surface in $\mathbb{P}^3(w_0, w_1, w_2, w_3)$ defined by the equation

$$x_0^2 = f(x_1, x_2, x_3) \quad \text{or} \quad x_0^2 x_i = f(x_1, x_2, x_3)$$

where σ acts on it by $\sigma(x_0) = -x_0$. We assume that $Q = (w_0, w_1, w_2, w_3)$ is normalized. The algorithm described in this section also works for more general $K3$ surfaces in $\mathbb{P}^3(Q)$ with non-symplectic involution.

Since S_0 is singular, S is chosen to be the minimal resolution of S_0 as in

$$\begin{tikzcd}
\mathbb{P}^3(w_0, w_1, w_2, w_3) & \tilde{\mathbb{P}^3}(w_0, w_1, w_2, w_3) \arrow[l] \\
\cup & \cup \\
S_0 & S \arrow[l]
\end{tikzcd}$$

where $\tilde{\mathbb{P}^3}(w_0, w_1, w_2, w_3)$ is a partial resolution of $\mathbb{P}^3(w_0, w_1, w_2, w_3)$ so that S is non-singular. The curve C_g is the strict transform of the curve defined by $x_0 = 0$ on S_0 which is isomorphic to the curve $f(x_1, x_2, x_3) = 0$ in $\mathbb{P}^2(w_1, w_2, w_3)$. The rational curves L_i are among those defined by letting either x_1, x_2 or x_3 be zero, or from the exceptional divisors arising in the desingularization. The procedure is as follows.

(i) The curve $\{x_0 = 0\}$ on S_0 is fixed by σ. Since $f(x_1, x_2, x_3) = 0$ is quasi-smooth (and hence smooth) in $\mathbb{P}^2(w_1, w_2, w_3)$, its strict transform C_g is also fixed by σ. The genus g can be calculated from $d := \deg f$ and weight (w_1, w_2, w_3) once it is normalized (see examples below). For instance, one can use the formula:

$$g = \frac{1}{2}\left(\frac{d^2}{w_1w_2w_3} - d\sum_{i>j}\frac{\gcd(w_i, w_j)}{w_iw_j} + \sum_{i=1}^{3}\frac{\gcd(d, w_i)}{w_i} - 1\right).$$

(ii) The locus $\{x_1 = 0\}$, $\{x_2 = 0\}$ or $\{x_3 = 0\}$ may be fixed by σ depending on (w_0, w_1, w_2, w_3). For instance, consider the case where w_0 is odd. If w_2 and w_3 are even (and w_3 is necessarily odd), then the locus $\{x_3 = 0\}$ is fixed by σ. In this case, the strict transform of the locus $\{x_3 = 0\}$ is ramified under the quotient $S \to S/\sigma$. It can be shown that this locus is a rational curve on S and contributes to a curve of L_i's in Lemma 5.1.

(iii) The rest of the curves in L_is are from exceptional divisors in the minimal resolution $S_0 \leftarrow S$. To find the divisors where σ acts as identity, we look carefully at the σ-action around the singularities of S_0 fixed by σ. Every singularity $P = (x_0 : x_1 : x_2 : x_3)$ on S_0 has at least two coordinates zero. Hence P is fixed by σ if and only if

- it has three zero coordinates, or
- it has exactly two zero coordinates with $x_0 = 0$, or
- it has exactly two zero coordinates and two non-zero coordinates x_0, x_i, and if $d = \gcd(w_0, w_i) \geq 2$, then w_0/d is odd and w_i/d is even.

Combining information obtained in (ii) and (iii), we can calculate the number k and hence the invariants r and a by the formula given in Lemma 5.1).

Computation of g and k for surfaces $x_0^2 = f(x_1, x_2, x_3)$

We consider surfaces $x_0^2 = f(x_1, x_2, x_3)$ in detail. Since the minimal resolution S is a $K3$ surface, P is a cyclic quotient singularity of type $A_{n+1,n}$ (or simply A_n) for some positive integer n. We obtain n exceptional divisors by resolving P. As P is fixed by σ, σ sends every exceptional divisor to itself. However, the σ action on each divisor is either $+1$ or -1, which depends on the singularity. The detail is explained in the following lemmas.

Lemma 5.2. *Let $S_0 :\ x_0^2 = f(x_1, x_2, x_3)$ be one of the 48 $K3$ surfaces defined in* Proposition 4.1. *If $w_1 \geq 2$ and $P = (0, 1, 0, 0)$ is on S_0, then P is a cyclic quotient singularity of type A_{w_1, w_1-1} $(= A_{w_1-1})$. Among the exceptional divisors arising from P, ramified divisors appear alternately and there are $\left[\dfrac{w_1-1}{2}\right]$ of them, where $[x]$ denotes the integer part of x. The same assertion holds for singularities $(0, 0, 1, 0)$ and $(0, 0, 0, 1)$.*

Proof. When $P = (0, 1, 0, 0)$ is a singularity, depending on $f(x_1, x_2, x_3)$, (x_0, x_2) or (x_0, x_3) gives a pair of local coordinates around P. Assume that (x_0, x_2) is a coordinate system. Since P is of type A_{w_1,w_1-1}, the μ_{w_1}-action above P can be written as

$$(x_0, x_2) \longmapsto (\zeta^{w_1-1}x_0, \zeta x_2)$$

with $\zeta \in \mu_{w_1}$. By the quotient map $S_0 \longrightarrow S_0/\sigma$, P is mapped to $(0,1,0,0) \in S_0/\sigma$ and the group action around this point is

$$(y_0, x_2) \longmapsto (\zeta^{2(w_1-1)}y_0, \zeta x_2)$$

with $y_0 = x_0^2$. Hence it is of type $A_{w_1,2(w_1-1)}$ (or precisely, A_{w_1,w_1-2}). In order to see how σ acts on the exceptional divisors, consider the continued fractional expansions

$$\frac{w_1}{w_1-1} = 2 - \cfrac{1}{2 - \cfrac{1}{2 - \cfrac{1}{2 - \cfrac{1}{\cdots}}}} \quad \text{and} \quad \frac{w_1}{2(w_1-1)} = 1 - \cfrac{1}{4 - \cfrac{1}{1 - \cfrac{1}{4 - \cfrac{1}{\cdots}}}}$$

This shows that if $E_1 + E_2 + E_3 + \cdots$ are exceptional divisor arising from P, then

$$\sigma(E_i)^2 = \begin{cases} -1 & \text{if } i \text{ is odd} \\ -4 & \text{if } i \text{ is even.} \end{cases}$$

By the projection formula, we see that σ acts as identity (resp. -1) on E_i for even i (resp. odd i). Therefore the exceptional divisors E_i with even i are fixed by σ and there are $[\frac{w_1-1}{2}]$ such divisors. $\square$

Next, we discuss singularities with exactly two zero coordinates, one of which is $x_0 = 0$.

Lemma 5.3. *Let $S_0 : x_0^2 = f(x_1, x_2, x_3)$ be one of the 48 $K3$ surfaces defined in* Proposition 4.1. *If $d := \gcd(w_1, w_2) \geq 2$, then $P = (0, x_1, x_2, 0)$ with $x_1x_2 \neq 0$ is a singularity of S_0 and fixed by σ. Among the exceptional divisors arising from P, there are $\left[\dfrac{d-1}{2}\right]$ ramified divisors and they appear alternately. The same assertion holds for singularities $(0, x_1, 0, x_3)$ and $(0, 0, x_2, x_3)$.*

Proof. When $P = (0, x_1, x_2, 0)$ is a singularity, (x_0, x_3) gives a local coordinate system around P; that is, P is a singularity of the quotient of an affine plane by the group action μ_d defined by

$$(x_0, x_3) \longmapsto (\zeta^{w_0}x_0, \zeta^{w_3}x_3)$$

with $\zeta \in \mu_d$. Since the weight is normalized, $\gcd(d, w_0) = \gcd(d, w_3) = 1$ and the μ_d action above can be written as

$$(x_0, x_3) \longmapsto (\zeta^{d-1}x_0, \zeta x_3).$$

(Note that P is a singularity of type $A_{d,d-1}$.) By the quotient map $S_0 \longrightarrow S_0/\sigma$, P is mapped to $(0, x_1, x_2, 0) \in S_0/\sigma$. The group action around this point is

$$(y_0, x_3) \longmapsto (\zeta^{2(d-1)}y_0, \zeta x_3)$$

with $y_0 = x_0^2$. Hence it is of type $A_{d,2(d-1)}$ (or precisely, $A_{d,d-2}$). As in Lemma 5.2, if $E_1 + E_2 + E_3 + \cdots$ are exceptional divisor arising from P, then only the exceptional divisors E_i with even i are fixed by σ. Therefore ramified exceptional divisors appear alternately and there are $[\frac{d-1}{2}]$ such divisors. $\square$

Lastly, we look at the singularity with two zero coordinates and $x_0 \neq 0$.

Lemma 5.4. *Let $S_0 : \ x_0^2 = f(x_1, x_2, x_3)$ be one of the* 48 *$K3$ surfaces defined in* Proposition 4.1. *If $d := \gcd(w_0, w_1) \geq 2$, then $P = (x_0, x_1, 0, 0)$ with $x_0x_1 \neq 0$ is a singularity of S_0. It is fixed by σ if and only if w_0/d is odd and w_1/d is even. Let $E_1 + \cdots + E_{d-1}$ denote the exceptional divisors arising from P. Then σ acts on E_i as identity (resp. by -1) if i is odd (resp. even). There are in total $\left[\dfrac{d}{2}\right]$ divisors with odd i. The same assertion holds for singularities $(x_0, 0, x_2, 0)$ and $(x_0, 0, 0, x_3)$.*

Proof. Since S_0 is quasi-smooth, one knows that $P = (x_0, x_1, 0, 0)$ with $x_0x_1 \neq 0$ is a singularity if and only if $d = \gcd(w_0, w_1) \geq 2$. Let $w_0 = du_0$ and $w_1 = du_1$ with $\gcd(u_0, u_1) = 1$. To see if P is fixed by σ, there are three cases to consider: $(u_0, u_1) = (\text{even}, \text{odd}), (\text{odd}, \text{odd}), (\text{odd}, \text{even})$. Suppose that there is t such that $t^{w_0} = -1$ and $t^{w_1} = 1$. If u_0 is even and u_1 is odd, then $t^{u_0u_1d} = (t^{u_0d})^{u_1} = (-1)^{u_1} = -1$ and $t^{u_0u_1d} = (t^{u_1d})^{u_0} = 1$, which is absurd. By the same reason as above, $(u_0, u_1) = (\text{odd, odd})$ cannot happen either. If u_0 is odd and u_1 is even, then let ζ be a primitive $2d$-th root of unity. We have $\zeta^{w_0} = (\zeta^d)^{u_0} = (-1)^{u_0} = -1$ and $\zeta^{w_1} = (\zeta^d)^{u_1} = (-1)^{u_1} = 1$. Hence there does exist t satisfying $t^{w_0} = -1$ and $t^{w_1} = 1$, and P is fixed by σ in this case.

Choose (x_2, x_3) as a local coordinate system. P is isomorphic to the singularity of the quotient by the group action μ_d defined by

$$(x_2, x_3) \longmapsto (\zeta^{w_2}x_2, \zeta^{w_3}x_3)$$

where ζ runs through μ_d. Since the weight is normalized, $\gcd(d, w_2) = \gcd(d, w_3) = 1$ and the μ_d action above can be written as

$$(x_2, x_3) \longmapsto (\zeta^{d-1}x_2, \zeta x_3).$$

P is mapped to $(x_0, x_1, 0, 0) \in S_0/\sigma$ and the group action around this point is

$$(x_2, x_3) \longmapsto (\zeta^{d-1}x_2, \zeta x_3)$$

as above. But, since $\gcd(2w_0, w_1) = \gcd(2du_0, du_1) = 2d\gcd(u_0, u_1/2) = 2d$, ζ now runs through μ_{2d}. Hence this singularity on S_0/σ is of type $A_{2d,d-1}$.

Consider two continued fractions

$$\frac{d}{d-1} = 2 - \cfrac{1}{2 - \cfrac{1}{2 - \cfrac{1}{2 - \cfrac{1}{\cdots}}}} \quad \text{and} \quad \frac{2d}{d-1} = 4 - \cfrac{1}{1 - \cfrac{1}{4 - \cfrac{1}{1 - \cfrac{1}{\cdots}}}}$$

This shows that if $E_1 + E_2 + E_3 + \cdots$ are exceptional divisor arising from P, then only the exceptional divisors E_i with odd i are fixed by σ. Therefore ramified exceptional divisors appear alternately and there are $[\frac{d}{2}]$ such divisors. □

Recall that r is the rank of $\mathrm{Pic}\,(S)^\sigma$. If we have good knowledge of $\mathrm{Pic}\,(S)$ and the σ action on it, then we can compute r without knowing S^σ. The following theorem shows that we can in fact find a closed formula for r by taking this approach.

Theorem 5.5. *Let (S, σ) be one of the* 48 *$K3$ surfaces defined in* Proposition 4.1 *as the minimal resolution of a hypersurface*

$$S_0 : \ x_0^2 = f(x_1, x_2, x_3) \quad \subset \mathbb{P}^3(Q)$$

where $Q = (w_0, w_1, w_2, w_3)$ *and* f *is a homogeneous polynomial of degree* $w_0 + w_1 + w_2 + w_3$. *Recall* $r = \operatorname{rank} \operatorname{Pic}(S)^\sigma$. *Let* $r(Q)$ *denote the number of exceptional divisors in the resolution* $S \longrightarrow S_0$. *Assume that* $\operatorname{rank} \operatorname{Pic}(S_0)^\sigma = 1$.

(a) *If* w_0 *is odd, then there exists at most one odd weight* w_i *such that* $\gcd(w_0, w_i) \geq 2$ *and in such a case,* $\gcd(w_0, w_i) = w_i$. *We have*

$$r = \begin{cases} r(Q) - w_i + 2 & \textit{if } \gcd(w_0, w_i) \geq 2 \textit{ for some odd weight } w_i \ (1 \leq i \leq 3) \\ r(Q) + 1 & \textit{otherwise.} \end{cases}$$

(b) *If* w_0 *is even, then let* $d_i = \gcd(w_0, w_i)$. *We have*

$$r = r(Q) + 1 - \sum_{i=1}^{3} (d_i - 1)\left(\frac{2d_i}{w_i} - 1\right).$$

Proof. Since $\operatorname{Pic}(S_0) \cong \mathbb{Z}$, $\operatorname{Pic}(S_0) \otimes \mathbb{Q}$ is generated by the hyperplane section $\{x_0 = 0\}$, which is fixed by σ. Its strict transform on S is also fixed by σ and gives a one-dimensional subspace of $\operatorname{Pic}(S)^\sigma \otimes \mathbb{Q}$. The rest of it is generated by exceptional divisors.

Possible singularities for S_0 are either of the form $(0 : x_1 : x_2 : x_3)$ with one or two zero coordinates or of the form $(x_0 : x_1 : x_2 : x_3)$ with $x_0 \neq 0$ and exactly two zero coordinates. Since the points with $x_0 = 0$ are fixed by σ, every exceptional divisor E arising from a singularity $(0 : x_1 : x_2 : x_3)$ satisfies $\sigma(E) = E$. (σ acts on E as ± 1.) Such exceptional divisors form a part of a basis for $\operatorname{Pic}(S)^\sigma$.

Consider a singularity $(x_0 : x_1 : x_2 : x_3)$ with $x_0 x_i \neq 0$, $\gcd(w_0, w_i) \geq 2$ and other coordinates zero. It is fixed by σ if and only if $t^{w_0} = -1$ and $t^{w_i} = 1$ for some t. As $x_0 x_i \neq 0$ and other coordinates are zero, $f(x_1, x_2, x_3)$ contains a monomial solely in x_i. Hence $w_i \mid 2w_0$, where $2w_0 = \deg f$. Now we divide the proof according as the parity of w_0.

(a) Assume that w_0 is odd. Then the normality of weight Q and the equality $w_0 = w_1 + w_2 + w_3$ imply that there is exactly one odd w_i for $1 \leq i \leq 3$. For simplicity, assume that w_1 is odd and w_2 and w_3 are even.

(i) If $\gcd(w_0, w_1) \geq 2$, then $(x_0 : x_1 : 0 : 0)$ are singular points. Since $1 = (t^{w_1})^{w_0} = (t^{w_0})^{w_1} = -1$, none of them is fixed by σ. But, if we consider a σ-conjugate pair, it is invariant under σ. There are two singularities of the form $(x_0 : x_1 : 0 : 0)$, each of which is of type $A_{w_1 - 1}$. (As $w_1 \mid 2w_0$ and w_1 is odd, we have $\gcd(w_0, w_1) = w_1$.) Totally, there are $2(w_1 - 1)$ exceptional divisors arising from these singularities and $w_1 - 1$ conjugate pairs contribute to $r = \operatorname{rank} \operatorname{Pic}(S)^\sigma$.

If $\gcd(w_0, w_1) = 1$, then $(x_0 : x_1 : 0 : 0)$ is not a singularity.

(ii) Consider the case where $\gcd(w_0, w_2) \geq 2$ and w_2 is even. $(x_0 : 0 : x_2 : 0)$ is a singularity. By letting $t = -1$, we see $t^{w_0} = -1$ and $t^{w_2} = 1$. Hence $(x_0 : 0 : x_2 : 0)$ is fixed by σ and so are the exceptional divisors arising from this point. This means that all exceptional divisors contribute to r. The same argument is valid for the singularities $(x_0 : 0 : 0 : x_3)$ with even w_3.

Therefore it follows from (i) and (ii) that $r = r(Q) + 1$ if there is no odd w_i with $\gcd(w_0, w_i) \geq 2$, and

$$r = r(Q) + 1 - (w_i - 1) = r(Q) - w_i + 2$$

if $\gcd(w_0, w_i) \geq 2$ for some odd weight w_i.

(b) Assume now that w_0 is even. Then the normality of weight Q and the equality $w_0 = w_1 + w_2 + w_3$ imply that there is exactly one even weight w_i for $1 \le i \le 3$. For simplicity, let w_1 be even, and w_2 and w_3 be odd.

(i) Consider the point $(x_0 : x_1 : 0 : 0)$ with w_1 even. Since $\gcd(w_0, w_1) \ge 2$, it is a singularity. We see $w_1 \mid 2w_0$.

If $w_1 \mid w_0$, then $d_1 = \gcd(w_0, w_1) = w_1$ and $\{(x_0 : x_1 : 0 : 0)\}$ consists of two points, each of which is a singularity of type A_{w_1-1}. There arise $2(w_1 - 1)$ exceptional divisors from them and $w_1 - 1$ conjugate pairs are fixed by σ. This means that the rank r is the number of exceptional divisors minus $w_1 - 1$.

If $w_1 \not| \, w_0$, then $\operatorname{lcm}(w_0, w_1) = 2w_0$ and $(x_0 : x_1 : 0 : 0)$ is a singularity of type A_{d_1-1}. This point is fixed by σ and so are the exceptional divisors arising from it.

In summary, the rank r is less than the number of exceptional divisors by

$$-(d_1 - 1)\left(\frac{2w_0}{\operatorname{lcm}(w_0, w_1)} - 1\right) = -(d_1 - 1)\left(\frac{2d_1}{w_1} - 1\right).$$

(ii) Consider the points $(x_0 : 0 : x_2 : 0)$ with w_2 odd. They are singularities if and only if $\gcd(w_0, w_2) \ge 2$. Here $w_2 \mid 2w_0$ implies $w_2 \mid w_0$ and $d_2 = \gcd(w_0, w_2) = w_2$. Such singularities are of type A_{d_2-1}. The multiplicity of $(x_0 : 0 : x_2 : 0)$ is $2w_0/\operatorname{lcm}(w_0, w_2) = 2$ and they are σ-conjugate. Among the $2(w_2 - 1)$ exceptional divisors, $w_2 - 1$ conjugate pairs are fixed by σ. Hence rank r is less than the number of exceptional divisors by

$$-(d_2 - 1)\left(\frac{2d_2}{w_2} - 1\right).$$

The same argument holds for the points $(x_0 : 0 : 0 : x_3)$ with odd w_3.

Therefore the asserted formula of (b) follows from (i) and (ii). □

Remark 5.1. For the $K3$ surfaces defined by $x_0^2 = f(x_1, x_2, x_3)$, σ acts as identity on the strict transform of the hyperplane section $\{x_0 = 0\}$. On some other hyperplane sections, σ may act as -1. But, if we regard them as divisors, they are invariant under the action of σ.

Example 5.6. (1) For $Q = (7, 3, 2, 2)$, we find that w_0 is odd, $r(Q) = 9$ and no odd weight w_i with $\gcd(w_0, w_i) \ge 2$. Hence $r = 9 + 1 = 10$.

(2) For $Q = (15, 10, 3, 2)$, we find that w_0 is odd, $r(Q) = 11$ and $\gcd(15, 3) = 3$. Hence $r = 11 - 3 + 2 = 10$.

(3) For $Q = (8, 4, 3, 1)$, we find that w_0 is even, $r(Q) = 8$ and $d_1 = 4$. Hence $r = 8 + 1 - (4 - 1)(2 \cdot 4/4 - 1) = 6$.

(4) For $Q = (10, 5, 3, 2)$, we find that w_0 is even, $r(Q) = 12$, $d_1 = 5$ and $d_3 = 2$. Hence $r = 12 + 1 - (5 - 1)(2 \cdot 5/5 - 1) - (2 - 1)(2 \cdot 2/2 - 1) = 8$.

(5) For $Q = (24, 16, 5, 3)$, we find that w_0 is even, $r(Q) = 15$, $d_1 = 8$ and $d_3 = 3$. Hence $r = 15 + 1 - (8 - 1)(2 \cdot 8/16 - 1) - (3 - 1)(2 \cdot 3/3 - 1) = 14$.

Example 5.7. Take a look at the $K3$ surface #8 in Yonemura defined by the equation

$$S_0 : \; x_0^2 = x_1^4 + x_2^6 + x_3^{12} \quad \subset \mathbb{P}^3(6, 3, 2, 1).$$

The involution is defined by $\sigma(x_0) = -x_0$.

We see that S_0 is quasi-smooth and the minimal resolution S is a $K3$ surface. S_0 has four singularities, P_1, P_1', P_2 and P_2' as follows:

Singularity	Type	Exceptional divisors
$P_1 := (1:1:0:0)$	$A_{3,2}$	$E_1 + E_2$
$P_1^{'} := (-1:1:0:0)$	$A_{3,2}$	$E_1^{'} + E_2^{'}$
$P_2 := (1:0:1:0)$	$A_{2,1}$	E_3
$P_2^{'} := (-1:0:1:0)$	$A_{2,1}$	$E_3^{'}$

No singularity is fixed by σ. There is a curve, $C^{'}$, fixed by σ, which is defined by $x = 0$. $C^{'}$ has genus 7. Its strict transform C $(= C_7)$ on S is ramified under σ. We have

$$S^\sigma = C_7.$$

Hence $g = 7$ and $k = 0$. This implies that $r = 11-7+0 = 4$ and $a = 11-7-0 = 4$.

The four divisors on S fixed by σ are, for instance, $E_1+E_1^{'}$, $E_2+E_2^{'}$, $E_3+E_3^{'}$ and C_7. But, by calculating the determinant of their intersection matrix, we see that they do not form a $\mathbb{Z}$-basis for $\mathrm{Pic}\,(S)^\sigma$. To find a right basis, we need to choose for C_7 the rational curve D defined by $x_3 = 0$. Note that when D is considered as a divisor, it is fixed by σ.

Since D is a rational curve on a $K3$ surface, $D^2 = -2$. As the curve $\{x_3 = 0\}$ on S_0 passes through every singularity,

$$D.(E_1 + E_1^{'}) = D.(E_3 + E_3^{'}) = 2, \quad D.(E_2 + E_2^{'}) = 0.$$

Hence the determinant of the intersection matrix for the new basis is

$$\begin{vmatrix} -4 & 2 & 0 & 2 \\ 2 & -4 & 0 & 0 \\ 0 & 0 & -4 & 2 \\ 2 & 0 & 2 & -2 \end{vmatrix} = -2^4$$

which agrees with the above calculation of $a = 4$.

It is not too difficult to find some subgroup of $\mathrm{Pic}\,(S_0)^\sigma$, but very difficult to describe $\mathrm{Pic}\,(S_0)^\sigma$ completely. Above calculations give us a clew to determine $\mathrm{rank}\,\mathrm{Pic}\,(S_0)^\sigma$ over $\mathbb{Q}$. In fact, let S be the minimal resolution of S_0 and let $\mathbb{E}$ denote the subgroup of $\mathrm{Pic}\,(S)$ generated by the exceptional divisors of the resolution. We have

$$\mathrm{Pic}\,(S)^\sigma \cong \mathrm{Pic}\,(S_0)^\sigma \oplus \mathbb{E}^\sigma.$$

Groups $\mathbb{E}$ and $\mathbb{E}^\sigma$ are easily describable and the rank of $\mathrm{Pic}\,(S)^\sigma$ may be computed from the fixed part S_0^σ by Lemma 5.1 as

$$\mathrm{rank}\,\mathrm{Pic}\,(S_0)^\sigma = \mathrm{rank}\,\mathrm{Pic}\,(S)^\sigma - \mathrm{rank}\,\mathbb{E}^\sigma = 11 + k - g - \mathrm{rank}\,\mathbb{E}^\sigma.$$

Hence if one already knows this number of divisors in $\mathrm{Pic}\,(S_0)^\sigma$, then these divisors form a basis for $\mathrm{Pic}\,(S_0)^\sigma$ over $\mathbb{Q}$.

Corollary 5.8. *Let (S, σ) be one of the 48 $K3$ surfaces with involution σ defined by a hypersurface of the form $x_0^2 = f(x_1, x_2, x_3)$ where σ acts by $\sigma(x_0) = -x_0$. Let $S^\sigma = C_g \cup L_1 \cup \ldots \cup L_k$ be the disjoint decomposition of S^σ, where C_g is a smooth genus g curve and $L_1, \cdots, L_k$ are rational curves.*

Suppose that f is defined by three monomials, so that S is of Delsarte type. Then the Jacobian variety $J(C_g)$ of C_g is also of CM type.

Proof. In this case C_g is defined by putting $x_0 = 0$. So C_g is defined by three monomials and is realized as a Fermat quotient. □

In the next section, we discuss another type (non-Borcea type) of our $K3$ surfaces. Putting an emphasis on the differences from the case $x_0^2 = f(x_1, x_2, x_3)$, we sketch the outline of our algorithm.

6. Computations of Nikulin's invariants for $K3$ surfaces of non-Borcea type

In this section, we compute r and a for $K3$ surfaces of non-Borcea type, namely for a quasi-smooth $K3$ surface S_0 in $\mathbb{P}^3(w_0, w_1, w_2, w_3)$ defined by the equation

$$x_0^2 x_i = f(x_1, x_2, x_3) \tag{6.1}$$

for some i $(= 1, 2, 3)$. Let S be the minimal resolution of S_0. Write C_g for the strict transform of the curve defined by $x_0 = 0$, which is isomorphic to the curve $f(x_1, x_2, x_3) = 0$ in $\mathbb{P}^2(w_1, w_2, w_3)$. We assume that $f(x_1, x_2, x_3) = 0$ is quasi-smooth (hence smooth) in $\mathbb{P}^2(w_1, w_2, w_3)$ after normalization of the weight.

Since $Q = (w_0, w_1, w_2, w_3)$ is normalized, every fixed point in S_0^σ must have at least one zero coordinate. There are four cases to consider.

(i) The curve $\{x_0 = 0\}$ on S_0 is fixed by σ. Since $f(x_1, x_2, x_3) = 0$ is quasi-smooth in $\mathbb{P}^2(w_1, w_2, w_3)$, its strict transform C_g is also fixed by σ. The genus g can be calculated from $\deg f$ and weight (w_1, w_2, w_3) as in the previous section.

The rational curves L_i of $S^\sigma = C_g \cup L_1 \cup \cdots \cup L_k$ are obtained by letting x_1, x_2 or x_3 be zero, or from the exceptional divisors arising in the desingularization.

(ii) As in the case of $K3$ surfaces of Borcea type , the one-dimensional locus $\{x_1 = 0\}$, $\{x_2 = 0\}$ or $\{x_3 = 0\}$ may be fixed by σ. In addition to them, there may be another one-dimensional locus fixed by σ; it has two zero coordinates, one of which is x_i of (6.1).

For instance, consider a surface $x_0^2 x_1 = f(x_1, x_2, x_3)$ with $f(0, 0, x_3) = 0$. If w_0 is odd and w_3 is even, then the locus $(x_0 : 0 : 0 : x_3)$ is a line and fixed by σ. Its strict transform on S is also fixed by σ, which gives one L_i of Lemma 5.1.

(iii) The rest of the curves in L_is are obtained from exceptional divisors in the resolution $S_0 \leftarrow S$. The singularities discussed in Lemmas 5.2, 5.3 and 5.4 also exist on surfaces (6.1) and by the same procedure as described there, we can find those divisors fixed identically by σ.

(iv) In addition to the singularities of (iii), we now have a singularity $(1 : 0 : 0 : 0)$ on the surface $x_0^2 x_i = f(x_1, x_2, x_3)$. The exceptional divisors arising from it and fixed by σ are determined as follows.

Lemma 6.1. *Let $S_0 :\ x_0^2 x_i = f(x_1, x_2, x_3)$ be one of the $K3$ surfaces obtained in* Theorem 4.2. *Then $P = (1, 0, 0, 0) \in S_0$ is a (cyclic quotient) singularity of type $A_{w_0, w_0 - 1}$. Let $E_1 + \cdots + E_{w_0 - 1}$ be the exceptional divisors on S arising from $P \in S_0$. On the quotient S_0/σ, we see that $\sigma(P)$ is a singularity of type $A_{2w_0, w_0 - 1}$ or $A_{w_0, 2(w_0 - 1)}$.*

(1) *If $\sigma(P)$ is of type $A_{2w_0, w_0 - 1}$, then σ acts on E_i as identity if and only if i is odd. There exist $\left[\dfrac{w_0}{2}\right]$ such divisors in total.*

(2) *If $\sigma(P)$ is of type $A_{w_0, 2(w_0 - 1)}$, then σ acts on E_i as identity if and only if i is even. There exist $\left[\dfrac{w_0 - 1}{2}\right]$ such divisors in total.*

The type of singularity at $\sigma(P)$ is determined as follows according to the parity of weights:

	w_0	w_i	w_j	w_k	*Type of* $\sigma(P)$
(a)	*even*	*even*	*odd*	*odd*	A_{2w_0,w_0-1}
(b)	*even*	*odd*	*odd*	*odd*	A_{2w_0,w_0-1}
(c)	*odd*	*odd*	*even*	*odd*	A_{2w_0,w_0-1}
$(c)'$	*odd*	*odd*	*odd*	*even*	A_{2w_0,w_0-1}
(d)	*odd*	*even*	*even*	*odd*	$A_{w_0,2(w_0-1)}$
$(d)'$	*odd*	*even*	*odd*	*even*	$A_{w_0,2(w_0-1)}$

Proof. Since $w_0 \geq 2$ and S is $K3$, $P = (1,0,0,0)$ is a cyclic quotient singularity of type A_{w_0,w_0-1}. From the equation $x_0^2 x_i = f(x_1, x_2, x_3)$, we can choose variables x_j and x_k, different from x_0 and x_i, as local parameters around P. The μ_{w_0}-action at P is then written as

$$(x_j, x_k) \longmapsto (\zeta^{w_j} x_j, \zeta^{w_k} x_k).$$

As $\deg f = 2w_0 + w_i$ and Q is a $K3$ weight, we have $2w_0 + w_i = w_0 + w_i + w_j + w_k$. This implies $w_0 = w_j + w_k$ and, because Q is normalized, $\gcd(w_0, w_j) = \gcd(w_0, w_k) = 1$. In particular, the congruence $w_j\alpha \equiv w_k \pmod{w_0}$ has solution $\alpha \equiv -1 \pmod{w_0}$. Now P is mapped to $(1,0,0,0) \in S_0/\sigma$ and the group action around this point is

$$(x_j, x_k) \longmapsto (\xi^{w_j} x_j, \xi^{w_k} x_k)$$

where ξ ranges over μ_{2w_0}. To find out the type of singularity at $\sigma(P)$, we divide the case according to the parity of weights. The cases (c) and $(c)'$, (d) and $(d)'$ are essentially the same, so we discuss cases (a), (b), (c) and (d).

In (a) and (b), w_0 is even. Since $w_0 = w_j + w_k$, both w_j and w_k are odd. Hence weight $(2w_0, w_i, w_j, w_k)$ is normalized and the singularity $(1,0,0,0) \in S_0/\sigma$ is of type A_{2w_0,w_0-1}. As in Lemma 5.4, we see that σ acts on E_i as identity if and only if i is odd, and there are $[w_0/2]$ such divisors.

In (c), w_0 is odd. Since $w_0 = w_j + w_k$, w_j and w_k have different parity. Because w_i is odd, the weight $(2w_0, w_i, w_j, w_k)$ is normalized. Hence the singularity $(1,0,0,0) \in S_0/\sigma$ is of type A_{2w_0,w_0-1}. As in Lemma 5.4, σ acts on E_i as identity if and only if i is odd, and there are $[w_0/2]$ such divisors.

In (d), w_0 is odd. Since $w_0 = w_j + w_k$, either w_j or w_k is even, say w_j is even. Because w_i is even in this case, $(2w_0, w_i, w_j, w_k)$ is not yet normalized. By normalization

$$\mathbb{P}^3(2w_0, w_i, w_j, w_k) \cong \mathbb{P}^3\Big(w_0, \frac{w_i}{2}, \frac{w_j}{2}, w_k\Big)$$

the group action around $(1,0,0,0) \in S_0/\sigma$ is now written as

$$(x_j, x_k) \longmapsto (\xi^{w_j/2} x_j, \xi^{w_k} x_k)$$

where ξ ranges over μ_{w_0}. As $\gcd(w_0, w_j/2) = \gcd(w_0, w_k) = 1$ and $w_0 = w_j + w_k$, we have the congruence

$$\frac{w_j}{2} 2(w_0 - 1) \equiv w_k \pmod{w_0}.$$

This shows that $(1,0,0,0) \in S_0/\sigma$ is of type $A_{w_0,2(w_0-1)}$ (to be more precise, type A_{w_0,w_0-2}). As in Lemma 5.3, σ acts on E_i as identity if and only if i is even, and there are $[(w_0-1)/2]$ such divisors. $\square$

We combine (ii), (iii) and (iv) to calculate the number of rational curves L_i. Then Lemma 5.1 gives the value for r and a. However, we are not able to obtain closed formula for r (similar to the formula obtained in Theorem 5.5) for these non-Borcea type $K3$ surfaces.

Example 6.2. Consider the $K3$ surface #60 in Yonemura. Dropping several monomials, we choose the equation

$$S_0 : \; x_0^2x_2 + x_1^3 + x_1x_2^3 + x_3^{18} = 0 \quad \subset \mathbb{P}^3(7,6,4,1).$$

The involution is defined by $\sigma(x_0) = -x_0$. We see that S_0 is quasi-smooth and the minimal resolution S is a $K3$ surface. S_0 has three singularities, P_1, P_2 and P_3 as follows:

Singularity	Type	Exceptional divisors
$P_1 := (1:0:0:0)$	$A_{7,6}$	$E_1 + E_2 + E_3 + E_4 + E_5 + E_6$
$P_2 := (0:0:1:0)$	$A_{4,3}$	$E_7 + E_8 + E_9$
$P_3 := (0:-1:1:0)$	$A_{2,1}$	E_{10}

Every singularity is fixed by σ, and E_2, E_4, E_6 and E_8 are ramified under σ (acting on the minimal resolution S).

There are two curves on S_0 fixed by σ, namely $C^{'}$ defined by $x_0 = 0$ and $L^{'}$ defined by $w = 0$. $C^{'}$ has genus 3 and $L^{'}$ is a projective line. Their strict transforms C $(= C_3)$ and L on S are ramified under σ. We have

$$S^\sigma = C_3 \cup E_2 \cup E_4 \cup E_6 \cup E_8 \cup L.$$

Hence $g = 3$ and $k = 5$. This implies $r = 13$ and $a = 3$.

Example 6.3. Consider the $K3$ surface #89 in Yonemura. Dropping several monomials, we choose the equation

$$S_0 : \; x_0^2x_3 + x_1^3x_2 + x_1x_2^4 + x_3^{11} = 0 \quad \subset \mathbb{P}^3(5,3,2,1).$$

The involution is defined by $\sigma(x_0) = -x_0$. We see that S_0 is quasi-smooth and the minimal resolution S is a $K3$ surface. S_0 has three singularities, P_1, P_2 and P_3 as follows:

Singularity	Type	Exceptional divisors
$P_1 := (1:0:0:0)$	$A_{5,4}$	$E_1 + E_2 + E_3 + E_4$
$P_2 := (0:1:0:0)$	$A_{3,2}$	$E_5 + E_6$
$P_3 := (0:0:1:0)$	$A_{2,1}$	E_7

Every singularity is fixed by σ, and E_2, E_4 and E_6 are ramified under σ (acting on the minimal resolution S).

There are two curves on S_0 fixed by σ, namely $C^{'}$ defined by $x_0 = 0$ and $L^{'}$ defined by $x_1 = x_3 = 0$. $C^{'}$ has genus 5 and $L^{'}$ is a projective line. Their strict transforms C $(= C_5)$ and L on S are ramified under σ. We have

$$S^\sigma = C_5 \cup E_2 \cup E_4 \cup E_6 \cup L.$$

Hence $g = 5$ and $k = 4$. This implies $r = 10$ and $a = 2$.

Corollary 6.4. *Let (S, σ) be one of the 86 $K3$ surfaces of Delsarte type with involution σ. Let C_g be the genus g curve in the fixed locus $S^\sigma = C_g \cup L_1 \cdots \cup L_k$ (where L_i are rational curves). Then C_g is of CM type in the sense that its Jacobian variety $J(C_g)$ is a CM abelian variety of dimension g.*

Proof. If σ acts as $\sigma(x_i) = -x_i$ on S, then C_g is defined by letting $x_i = 0$. It is a curve of Delstarte type and hence of CM type. $\square$

6.1. **Realization of Nikulin's invariants.** We briefly discuss how many Nikulin's triplets (r, a, δ) can be realized by our $K3$ surfaces. To realize as many triplets as possible, we introduce more involutions than those considered in previous sections.

First, we summarize the results of previous sections where σ acts on the variable x_0 (with the highest weight among x_i's).

Theorem 6.5. *Let* (S, σ) *be one of the* 92 $K3$ *surfaces in* Theorem 4.2 *with involution* $\sigma(x_0) = -x_0$. *Among the* 75 *triplets* (r, a, δ) *of Nikulin, at least* 29 *triplets are realized with such* $K3$ *surfaces. See* Tables 1 *to* 5 *and* 7 *of* Section 13 *for the list of defining equations of* S_0 *and the values for* (r, a)*; in most cases,* S_0 *is defined by*

$$x_0^2 = f(x_1, x_2, x_3) \quad \text{or} \quad x_0^2 x_i = f(x_1, x_2, x_3).$$

Remark 6.1. We say "at least 29" because we computed only the invariants r and a. If we are to find δ, we should have a $\mathbb{Z}$-basis for $\mathrm{Pic}\,(S)$ or calculate intersection numbers of divisors on S. We leave this as a future problem. Once δ is calculated, the number of realizable triplets may increase.

When Q is fixed and σ is defined by $\sigma(x_0) = -x_0$ on the surface $x_0^2 = f(x_1, x_2, x_3)$ or $x_0^2 x_i = f(x_1, x_2, x_3)$, no matter what quasi-smooth equation we choose for $f(x_1, x_2, x_3)$, the fixed locus S^σ is the same. Hence changes in equation $f(x_1, x_2, x_3)$ do not lead to any new pairs of r and a.

On the other hand, even with the same S_0, changing involution σ often changes the fixed locus S^σ, which has a possibility that we obtain a different pair of r and a. We use this approach by letting σ act on some variable x_i other than x_0.

Theorem 6.6. *Let* (S, σ) *be one of the* 92 $K3$ *surfaces in* Theorem 4.2 *having an involution* σ *on a variable* x_i *other than* x_0. *They are listed on* Table 9 *in* Section 13 *with values for* r *and* a. *Compared with the triplets obtained in* Theorem 6.5, *at least* 14 *more triplets are realized with such* σ *actions. Among the* 75 *triplets* (r, a, δ) *of Nikulin, the total number of triplets we realize is at least* 40. *The values for* r *and* a *of such* 40 *triplets are as follows:*

r	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
a	1	0 2	1			2 4	3 7	6 8	1 9	0 2 4 6 8	1 9	6 8	3 5 9	2 4 6	5 7	2 6	1 3 5	0 2 4	1 3	2

Proof. The σ-fixed locus S_0^σ can be determined by the same method as in Sections 5 and 6. In particular, the singularities of S_0 are independent of the choice of σ. We use Lemmas 5.2, 5.3, 5.4 and 6.1 to find which exceptional divisors are fixed by involution σ. $\square$

Remark 6.2. In some cases (say, Case #57), the σ-action on x_i $(i \neq 0)$ gives the same S^σ as the action by $\sigma(x_0) = -x_0$. Such cases are omitted from Table 9.

Remark 6.3. Hisanori Ohashi has communicated us an idea of even more different types of involutions on S_0. We thank him for the idea and plan to work on it in a subsequent paper.

7. Realization of Calabi–Yau threefolds as hypersurfaces over $\mathbb{Q}$

First we describe general construction of Calabi–Yau threefolds from product threefolds.

Let E be an elliptic curve over $\mathbb{Q}$ with involution ι. Let (S, σ) be a $K3$ surface with involution σ defined over $\mathbb{Q}$. Suppose that E and S have defining equations of the form

$$E : y^2 = g(x, z) \quad \text{with involution } \iota(y) = -y,$$

$$S : s^2 = f(u, v, w) \quad \text{with involution } \sigma(s) = -s$$

where $g(x, z)$ is a homogeneous cubic polynomial over $\mathbb{Q}$ in $\mathbb{P}^1(1, 1)$, and $f(u, v, w)$ is a homogeneous polynomial over $\mathbb{Q}$ in $\mathbb{P}^2(q_1, q_2, q_3)$ with some weight (q_1, q_2, q_3) giving rise to a $K3$ surface S. Then the product threefold $E \times S$ is defined by

$$(sy)^2 = g(x, z)f(u, v, w)$$

with involution $\iota \times \sigma$ in some weighted projective 4-space. The singularities are determined by weights. Perform resolution of singularities, to get a crepant resolution X, which is a Calabi–Yau threefold. We will have a model for X defined over $\mathbb{Q}$.

Now we will consider more explicit construction of Calabi–Yau threefolds, using twist maps (see Goto–Kloosterman–Yui [9] for details).

Let $\mathbb{P}^2(k + 1, k, 1)$ be a weighted projective 2-space with weight$(k + 1, k, 1)$ of degree $2(k+1)$. Let $\mathbb{P}^3(w_0, w_1, w_2, w_3)$ be a weighted projective 3-space with weight (w_0, w_1, w_2, w_3) of degree $d := \sum_{i=0}^{3} w_i$. Define a *twist map*:

$$\mathbb{P}^2(k + 1, k, 1) \times \mathbb{P}^3(w_0, w_1, w_2, w_3)$$

$$\mapsto \mathbb{P}^4(kw_0, w_0, (k + 1)w_1, (k + 1)w_2, (k + 1)w_3)$$

into a weighted projective 4-space with degree $(k + 1)d$. This will produce many Calabi–Yau threefolds defined by hypersurfaces in weighted projective 4-spaces.

Take $k = 1$. Then we have an elliptic curve

$$E_2 : y_0^2 = y_1^4 + y_2^4 \subset \mathbb{P}^2(2, 1, 1),$$

and if we take $k = 2$, we obtain an elliptic curve

$$E_3 : y_0^2 = y_1^3 + y_2^6 \subset \mathbb{P}^2(3, 2, 1).$$

Both E_2 and E_3 have complex multiplication, by $\mathbb{Z}[\sqrt{-1}]$ and $\mathbb{Z}[\sqrt{-3}]$, respectively.

Proposition 7.1 (Borcea [4]). *Let $E_2 : y_0^2 = y_1^4 + y_2^4 \subset \mathbb{P}^2(2, 1, 1)$ and $E_3 : y_0^2 = y_1^3 + y_2^6 \subset \mathbb{P}^2(3, 2, 1)$ be elliptic curves. Let $S : x_0^2 = f(x_1, x_2, x_3) \subset \mathbb{P}^3(w_0, w_1, w_2, w_3)$ be one of the* 40 *$K3$ surfaces from* Tables 1 *and* 2*.*

(a) *Suppose that w_0 is odd, and that $w_0 = w_1 + w_2 + w_3$. Then there is a twist map*

$$\mathbb{P}^2(2, 1, 1) \times \mathbb{P}^3(w_0, w_1, w_2, w_3) \cdots \longrightarrow \mathbb{P}^4(w_0, w_0, 2w_1, 2w_2, 2w_3)$$

given by

$$(y_0 : y_1 : y_2) \times (x_0 : x_1 : x_2 : x_3) \mapsto (y_1(\frac{x_0}{y_0})^{1/2} : y_2(\frac{x_0}{y_0})^{1/2} : x_1 : x_2 : x_3)$$

The product $E_2 \times S$ *maps generically* $2:1$ *to the hypersurface of degree* $4w_0$ *of the form*

$$z_0^4 + z_1^4 = f(z_2, z_3, z_4)$$

where f *is a homogeneous polynomial over* $\mathbb{Q}$ *of degree* $4w_0$*. This is a singular model for a Calabi–Yau threefold* $E_2 \times S/\iota \times \sigma$ *in* $\mathbb{P}^4(w_0, w_0, 2w_1, 2w_2, 2w_3)$*.*

(b) *Suppose that* w_0 *is even but not divisible by* 3*, and that* $w_0 = w_1 + w_2 + w_3$*. Then there is a twist map*

$$\mathbb{P}^2(3,2,1) \times \mathbb{P}^3(w_0, w_1, w_2, w_3) \cdots \longrightarrow \mathbb{P}^4(2w_0, w_0, 3w_1, 3w_2, 3w_3)$$

given by

$$(y_0 : y_1 : y_2) \times (x_0 : x_1 : x_2 : x_3) \mapsto (y_1(\frac{x_0}{y_0})^{2/3} : y_2(\frac{x_0}{y_0})^{1/3} : x_1 : x_2 : x_3)$$

The product $E_3 \times S$ *maps generically* $2:1$ *to the hypersurface of degree* $6w_0$ *of the form*

$$z_0^3 + z_1^6 = f(z_2, z_3, z_4)$$

where f *is a homogeneous polynomial over* $\mathbb{Q}$ *of degree* $6w_0$*. This is a singular model for a Calabi–Yau threefold* $E_3 \times S/\iota \times \sigma$ *in* $\mathbb{P}^4(2w_0, w_0, 3w_1, 3w_2, 3w_3)$*.*

Problem 1. When w_0 is divisible by 6, describe the twist map explicitly and construct hypersurface defining equations for Calabi–Yau threefolds modifying twist maps.

Theorem 7.2. *There are in total* 40 *Calabi–Yau threefolds realized by quasi-smooth hypersurfaces over* $\mathbb{Q}$ *in weighted projective* 4 *spaces by the above construction. On the other hand, if* S *is one of the* 45 $K3$ *surfaces of* Tables 1 *to* 3 *other than* #90, #91, #93*, then the associated* 45 *Calabi–Yau threefolds are all of CM type.*

Proof. Recall that the space of harmonic forms on a product is the tensor product of the corresponding spaces of harmonic forms of the components. Therefore, the Hodge structure h_X of a Calabi–Yau threefold X is given by the tensor product of the Hodge structures h_E and h_S of the components E and S respectively. We know that $h_X = h_E \otimes h_S$ is of CM type if and only if h_E and h_S are of CM type. With our choice of E and S, $E = E_2$ or E_3 is of CM type, and S is of CM type. Therefore, X is of CM type. □

For the additional 41 pairs (S, ι) of $K3$ surfaces with involution σ of Theorem 4.3 (b) (see Tables 4, 5 and 6), the situation is slightly different from the above cases. Calabi–Yau threefolds X are birational to hypersurfaces over $\mathbb{Q}$, but they are not quasi-smooth. More precisely, we have the following result.

Theorem 7.3. *Let* (S, σ) *be one of the* 41 *pairs of* $K3$ *surfaces with involution* σ *of* Theorem 4.3 (b)*, which is not in the list of Borcea. Let* E *be an elliptic curve over* $\mathbb{Q}$ *with involution* ι *with or without CM. Take the product* $E \times S$ *and consider the quotient threefold* $E \times S/\iota \times \sigma$*. Resolving singularities, we obtain a smooth Calabi–Yau threefold* X *over* $\mathbb{Q}$*. Further,* X *is of CM type if and only if* E *is of CM type.*

About the realization of X *as a hypersurface, the following holds.*

(a) *If (S, ι) is one of the $K3$ surfaces listed in* Tables 4 *and* 5 *other than* #22 *and* #58*, then S is defined by $x_0^2 x_i + f(x_1, x_2, x_3) = 0$ for some $i \neq 0$ and X is birational to a (non-quasi-smooth) hypersurface over $\mathbb{Q}$ defined by*

$$\begin{cases} z_{i+1}(z_0^4 + z_1^4) + f(z_2, z_3, z_4) = 0 & \text{if } w_0 \text{ is odd and } E = E_2 \\ z_{i+1}(z_0^3 + z_1^6) + f(z_2, z_3, z_4) = 0 & \text{if } w_0 \text{ is even but not divisible by } 3 \text{ and } E = E_3. \end{cases}$$

(b) *Let (S, ι) be one of the $K3$ surfaces listed in* Table 6 *other than* #16*. If we choose $E = E_2$, then X is birational to the following (non-quasi-smooth) hypersurface over $\mathbb{Q}$:*

$$\begin{cases} (z_0^4 + z_1^4)^2 + z_2^3 + z_3^4 + z_4^6 = 0 & \text{in } \#2 \\ (z_0^4 + z_1^4)^2 + z_2^3 + z_2 z_3^3 + z_3 z_4^4 = 0 & \text{in } \#52 \\ z_3(z_0^4 + z_1^4)^2 + z_2^3 + z_2 z_4^3 + z_3^3 z_4 = 0 & \text{in } \#84. \end{cases}$$

Proof. Since the singular locus of $E \times S/\iota \times \sigma$ is defined over $\mathbb{Q}$, resolving singularities, we obtain a smooth Calabi–Yau threefold X over $\mathbb{Q}$. Here X is not necessarily of CM type. By the same argument as for proof of Theorem 7.2, X is of CM type if and only if each compoment, E and S, is of CM type. Since we already know that S is of CM type, X is of CM type if and only if E is a CM elliptic curve over $\mathbb{Q}$.

(a) If we choose an appropriate elliptic curve E, then the twist map of Proposition 7.1 works for $x_0^2 = -f(x_1, x_2, x_3)/x_i$. Depending on the parity of w_0, we obtain the equations as claimed.

(b) Since the variable associated with the involution σ carries an odd weight, we may choose $E = E_2$. Then the twist map of Proposition 7.1 (a) works for $x_0^2 = \sqrt{-f(x_1, x_2, x_3)}$ or $x_0^2 = \sqrt{-f(x_1, x_2, x_3)/x_i}$ and we obtain the equations as asserted. ☐

Remark 7.1. Note that $K3$ surfaces S realized by Yonemura in weighted projective 3-spaces are often singular. To have smooth $K3$ surfaces, we ought to consider minimal resolutions $\tilde{S}$. The involution σ is lifted to $\tilde{S}$ and we use $\tilde{S}$ to carry out the above construction of Calabi–Yau threefolds X. The procedure is shown as follows:

$$\begin{tikzcd} E \times S & E \times \tilde{S} \arrow[l] \arrow[d] & \\ & E \times \tilde{S}/\iota \times \sigma & X \arrow[l] \end{tikzcd}$$

8. Singularities and resolutions on Calabi–Yau threefolds

Let S be a $K3$ surface defined by a weighted hypersurface

$$S : x_0^2 = f(x_1, x_2, x_3) \subset \mathbb{P}^3(w_0, w_1, w_2, w_3)$$

of degree $\deg(f) = w_0 + w_1 + w_2 + w_3$. The involution σ is given by $\sigma(x_0) = -x_0$. The singularities on S are determined by the weight. Let $\tilde{S}$ be the minimal resolution of S. The involution σ is extended to $\tilde{S}$. Let $\tilde{S}^\sigma$ be the fixed part of $\tilde{S}$ by σ.

Let E be an elliptic curve

$$E_2 : y_0^2 = y_1^4 + y_2^4 \subset \mathbb{P}^2(2, 1, 1)$$

or

$$E_3 : y_0^2 = y_1^3 + y_2^6 \subset \mathbb{P}^2(3, 2, 1).$$

The involution ι is given by $\iota(y_0) = -y_0$. The fixed part E^ι consists of four points

$$E^\iota = \{P_1, P_2, P_3, P_4\}.$$

We use for E either E_2 if w_0 is even, or E_3 if w_0 is odd. Take the quotient threefold $E \times S$.

Let X be a smooth resolution of $E \times \tilde{S}/\iota \times \sigma$. Then the singular loci $\{P_i\} \times \tilde{S}^\sigma$ are determined from the weight of S and the singularity data of the ambient space.

Here are examples.

Example 8.1. Let

$$S : x_0^2 = x_1^5 + x_2^5 + x_3^{10} \subset \mathbb{P}^3(5, 2, 2, 1).$$

(This #6 in Yonemura =#2 in Borcea.) Let

$$E_2 : y_0^2 = y_1^4 + y_2^4 \subset \mathbb{P}^2(2, 1, 1).$$

- Then S is singular $K3$ surface and the singular locus is:

$$\mathrm{Sing}(S) = \{(0 : x_1 : x_2 : 0) \,|\, x_1^5 + x_2^5 = 0\} = \{Q_1, Q_2, Q_3, Q_4, Q_5\}$$

where every Q_i is a cyclic quotient singularity of type A_1.

- Let $\bar{C}$ be a curve on S defined by $x_0 = 0$, that is,

$$\bar{C} = \{\, x_0 = 0\} : x_1^5 + x_2^5 + x_3^{10} = 0 \subset \mathbb{P}^2(2, 2, 1).$$

Since $\mathbb{P}^2(2, 2, 1) \simeq \mathbb{P}^2(1, 1, 1)$, $\bar{C}$ is identified with

$$\bar{C} : x_1^5 + x_2^5 + x_3^5 = 0 \subset \mathbb{P}^2$$

which is a smooth curve of genus 6.

- Let $\bar{L}$ be a curve on S defined by $x_3 = 0$, that is,

$$\bar{L} = \{\, x_3 = 0\,\} : x_0^2 = x_1^5 + x_2^5 \subset \mathbb{P}^2(5, 2, 2).$$

Since $\mathbb{P}^2(5, 2, 2) \simeq \mathbb{P}^2(5, 1, 1)$, $\bar{L}$ is identified with

$$\bar{L} : x_0 = x_1^5 + x_2^5 \subset \mathbb{P}^2(5, 1, 1)$$

and hence $\bar{L}$ is a rational curve.

- We see that

$$\bar{C} \cap \bar{L} = \{Q_1, Q_2, Q_2, Q_4, Q_5\}.$$

- Let $\tilde{S}$ be the minimal resolution of S. The involution σ is lifted to $\tilde{S}$. Let C_6 and L_1 be the respective strict transforms of $\bar{C}$ and $\bar{L}$ to the minimal resolution $\tilde{S}$. Let $\mathcal{E}_1, \cdots, \mathcal{E}_5$ be the exceptional divisors on $\tilde{S}$ arising from singularities Q_i, $(i = 1, \cdots, 5)$, respectively.

Then C_6 and L_1 are fixed by σ, but not the exceptional divisors $\mathcal{E}_i$ for any $i \in \{1, 2, \cdots, 5\}$. Hence

$$\tilde{S}^\sigma = C_6 \cup L_1$$

and we see $g = 6$ and $k = 1$ (so $r = 6$, $a = 4$) in the Voisin notation. The resolution picture is given in Figure 2, where the curves in boldface are fixed by σ.

Proposition 8.2. *Let X be a crepant resolution of the quotient threefold $E_2 \times \tilde{S}/\iota \times \sigma$ of* Example 8.1. *Then X is a Calabi–Yau threefold X corresponding to the triplet* $(6, 4, 0)$*, and its exceptional divisors are four copies of the ruled surfaces*

$$(C \times \mathbb{P}^1) \cup (L \times \mathbb{P}^1).$$


C_6
L_1
$\mathcal{E}_1$ $\mathcal{E}_2$ $\mathcal{E}_3$ $\mathcal{E}_4$ $\mathcal{E}_5$

Figure 2. Exceptional divisors and the fixed locus



Figure 3. Exceptional divisors and the fixed locus

Example 8.3. Consider the surface

$$S : x_0^2 = x_1^3x_3 + x_2^7 + x_3^{28} \subset \mathbb{P}^3(14, 9, 4, 1).$$

This is #45 in Yonemua=#36 in Borcea. Let

$$E_3 : y_0^2 = y_1^3 + y_2^6 \subset \mathbb{P}^2(3, 2, 1).$$

- The surface S is a singular $K3$ surface. There are two singular points:

$$Q := (0 : 1 : 0 : 0) \quad \text{of type } A_{9,8},$$

and

$$R := (1 : 0 : 1 : 0) \quad \text{of type } A_{2,1}.$$

- Let $\bar{C}$ be the curve on S defined by $x_0 = 0$

$$\bar{C} = \{x_0 = 0\} : x_1^3x_3 + x_2^7 + x_3^{28} = 0 \subset \mathbb{P}^2(9, 4, 1).$$

Then $\bar{C}$ is a quasi-smooth curve with singularity Q.

- No other curves defined by $x_i = 0$ $(i \neq 0)$ are fixed by the involution σ.
- Let $\tilde{S}$ be the minimal resolution of S. Then $\tilde{S}$ is a smooth $K3$ surface and the involution σ is lifted to $\tilde{S}$. Let C_6 be the strict transform of $\bar{C}$ to $\tilde{S}$; it has genus 6. Let $\mathcal{E}_1, \cdots, \mathcal{E}_8$ be exceptional divisors arising from singularity Q. Let $\mathcal{E}_9$ be the exceptional divisor arising from R. Then $\mathcal{E}_{2i}\,(i = 1, 2, 3, 4)$ and $\mathcal{E}_9$ are fixed by σ but not others.

Put $L_i := \mathcal{E}_{2i}\,(i = 1, 2, 3, 4)$ and $L_5 := \mathcal{E}_9$. Then

$$\tilde{S}^\sigma = C \cup L_1 \cup \cdots \cup L_5.$$

So $g = 6$ and $k = 5$ (so $r = 10, a = 0$) in the Voisin notation. The resolution picture is given in Figure 3, where the curves in boldface are fixed by σ.

- The quotient threefold $E_3 \times \tilde{S}/\iota \times \sigma$ has singularities along $\{P_i\} \times \tilde{S}^\sigma$ where $E_3^\iota = \{P_1, P_2, P_3, P_4\}$.

Summarizing the above, we have

Proposition 8.4. *A crepant resolution X of the quotient threefold $E_3 \times \tilde{S}/\iota \times \sigma$ of* Example 8.3 *is a Calabi–Yau threefold corresponding to the triplet* $(10, 0, 0)$*, and its exceptional divisors are four copies of*

$$(C \times \mathbb{P}^1) \cup (L_1 \times \mathbb{P}^1) \cup \cdots \cup (L_5 \times \mathbb{P}^1).$$

Example 8.5. Consider the surface

$$S : x_0^2 = x_1^3 + x_2^{10} + x_2^{15} \subset \mathbb{P}^3(15, 10, 3, 2).$$

This is #11 in Yonemura =#18 in Borcea. Let

$$E_2 : y_0^2 = y_1^4 + y_2^4 \subset \mathbb{P}^2(2, 1, 1).$$

- S is a singular $K3$ surface and singularities are

$$Q_1, Q_2, Q_3 = (0 : x_1 : 0 : x_3) \quad \text{of type } A_{2,1},$$

$$R := (x_0 : x_1 : 0 : 0) \quad \text{of type } A_{5,4},$$

and

$$T_1, T_2 := (x_0 : 0 : x_2 : 0) \quad \text{of type } A_{3,2}.$$

- Let $\bar{C}$ be the curve on S defined by $x_0 = 0$:

$$\bar{C} = \{x_0 = 0\} : x_1^3 + x_2^{10} + x_3^{15} = 0 \subset \mathbb{P}^2(10, 3, 2).$$

Via the isomorphism $\mathbb{P}^2(10, 3, 2) \simeq \mathbb{P}^2(5, 3, 1)$, $\bar{C}$ is identified with

$$\bar{C} : x_1^3 + x_2^5 + x_3^{15} = 0 \subset \mathbb{P}^2(5, 3, 1).$$

- Let $\bar{L}$ be the curve on S defined by $x_2 = 0$:

$$\bar{L} = \{x_2 = 0\} : x_0^2 = x_1^3 + x_3^{15} \subset \mathbb{P}^2(15, 10, 2).$$

Via the isomorphism $\mathbb{P}^2(15, 10, 2) \simeq \mathbb{P}^2(3, 1, 1)$, $\bar{L}$ is identified with

$$\bar{L} : x_0 = x_1^3 + x_3^3 \subset \mathbb{P}^2(3, 1, 1).$$

- $\bar{C}$ has genus 4 and $\bar{L}$ is rational with

$$\bar{C} \cap \bar{L} = \{Q_1, Q_2, Q_3\} \quad \text{and} \quad R \in \bar{L}.$$

- Let $\tilde{S}$ be the minimal resolution of S. The involution σ is lifted to $\tilde{S}$. Let C and L_1 be the strict transforms of $\bar{C}$ and $\bar{L}$ on S, respectively. Let $\mathcal{E}_i$ $(i = 1, 2, 3)$ be the exceptional divisors arising from singularities Q_i $(i = 1, 2, 3)$, $\mathcal{E}_{4+j}$ $(j = 0, 1, 2, 3)$ be the exceptional divisors arising from R, and $\mathcal{E}_{8+t}$ $(t = 0, 1, 2, 3)$ be the exceptional divisors arising from singularities T_1, T_2.
- $C, L_1, \mathcal{E}_5 =: L_2$ and $\mathcal{E}_7 =: L_3$ are fixed by σ, but all others are not. Hence

$$\tilde{S}^\sigma = C \cup L_1 \cup L_2 \cup L_3.$$

So $g = 4$ and $k = 3$ (so $r = 10, a = 4$) in the Voisin notation. The resolution picture is given in Figure 4, where the curves in boldface are fixed by σ.

- The quotient threefold $E_2 \times \tilde{S}/\iota \times \sigma$ has singularities $\{P_i\} \times \tilde{S}^\sigma$ $(i = 1, 2, 3, 4)$ where $E_2^\iota = \{P_1, P_2, P_3, P_4\}$.

Proposition 8.6. *A crepant resolution X of the quotient threefold $E_2 \times \tilde{S}/\iota \times \sigma$ of* Example 8.5 *is a Calabi–Yau threefold corresponding to the triplet* $(10, 4, 0)$*, and its exceptional divisors are four copies of ruled surfaces:*

$$(C \times \mathbb{P}^1) \cup (L_1 \times \mathbb{P}^1) \cup (L_2 \times \mathbb{P}^1) \cup (L_3 \times \mathbb{P}^1).$$



FIGURE 4. Exceptional divisors and the fixed locus

9. THE AUTOMORPHY OF CALABI–YAU THREEFOLDS OVER $\mathbb{Q}$

Let X be one of our Calabi–Yau varieties (elliptic curves, $K3$ surfaces, or Calabi–Yau threefolds). Let d be its dimension $d = 1, 2$, or 3. We will analyze the Galois representation associated to the middle dimensional ℓ-adic étale cohomology group $H^d_{et}(\bar{X}, \mathbb{Q}_\ell)$, where ℓ is a prime.

We say that p is a *good* prime if X has a model with good reduction X_p at p. Then X_p is a smooth Calabi–Yau variety over $\mathbb{F}_p$. The absolute Galois group $\mathrm{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ acts on $\bar{X} := X \otimes_{\mathbb{Q}} \bar{\mathbb{Q}}$, and one has an induced action on the cohomology group $H^i_{et}(\bar{X}, \mathbb{Q}_\ell)$ $(0 \leq i \leq 2d)$ of X, where ℓ is a prime different from p. This defines a continuous ℓ-adic representation $\rho : \mathrm{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \to GL_r(\mathbb{Q}_\ell)$ of some finite rank r. For a good prime p the structure of this Galois representation can be studied by passing to the reduction X_p. The Frobenius morphism Frob_p induces a $\mathbb{Q}_\ell$-linear map $\rho(\mathrm{Frob}_p)$ on $H^i_{et}(\bar{X}, \mathbb{Q}_\ell)$ $(i, 0 \leq i \leq 2d)$. Let

$$P^i_p(X, \rho, t) := \det(1 - \rho(\mathrm{Frob}_p)\, t, \,|\, H^i_{et}(\bar{X}, \mathbb{Q}_\ell))$$

be the characteristic polynomial of $\rho(\mathrm{Frob}_p)$, where t is an indeterminate. Then it is known that $P^i_p(X, \rho, t) \in 1 + \mathbb{Z}[t]$ has degree $B_i(X)$ (the i-th Betti number of X) and that its reciprocal roots are algebraic integers with complex absolute value $p^{i/2}$ (the Riemann Hypothesis for X_p). For each $i, 0 \leq i \leq 2d$, we define the i-th L-series by

$$L_i(X, s) := \prod_p \frac{1}{P^i_p(X, \rho, p^{-s})}.$$

In particular, the most important one is the middle-dimensional L-series $L_d(X, s)$.

We will now define a motivic L-series as follows.

Definition 9.1. Let X be a Calabi–Yau variety of dimension d defined over $\mathbb{Q}$. Consider the d-th étale ℓ-adic cohomology group $H^d_{et}(\bar{X}, \mathbb{Q}_\ell)$. Let $\mathcal{M}$ be a motive of X. Suppose that $\mathcal{M}$ is defined over $\mathbb{Q}$. Let $H^d_{et}(\mathcal{M}, \mathbb{Q}_\ell)$ be the étale ℓ-adic realization of $\mathcal{M}$. Let

$$P^d_p(\mathcal{M}, \rho, t) := \det(1 - \rho(\mathrm{Frob}_p)\, t \,|\, H^d_{et}(\bar{\mathcal{M}}, \mathbb{Q}_\ell)).$$

Then the motivic d-th L-series of $\mathcal{M}$ is defined by

$$L_d(\mathcal{M}, \rho, s) := \prod_p \frac{1}{P^d_p(\mathcal{M}, \rho, p^{-s})}.$$

If there is no danger of ambiguity, we may simply write $L(\mathcal{M}, \rho, s)$ dripping the subindex d.

Lemma 9.1. *(a) Let $d = 2$. Let S be one of the 86 $K3$ surfaces defined over $\mathbb{Q}$ in a weighted projective 3-space $\mathbb{P}^3(w_0, w_1, w_2, w_3)$ given in* Theorem 4.3. *Then the $K3$ motive $\mathcal{M}_S$ is the unique motive associated to the weight.*

(b) Let $d = 3$. Let X be one of our Calabi–Yau threefold defined over $\mathbb{Q}$ in a weighted projective 4-space $\mathbb{P}^4(w_0, w_1, w_2, w_3, w_4)$. Then the Calabi–Yau motive $\mathcal{M}_X$ is the unique motive associated to the weight.

For proof of Lemmas, the reader is referred to Gouvêa and Yui [10].

Definition 9.2. The unique motive $\mathcal{M}_Z$ with $h^{0,d}(\mathcal{M}_Z) = h^{d,0}(\mathcal{M}_Z) = 1$ is called the $K3$ *motive* if $d = 2$, and the *Calabi–Yau motive* if $d = 3$.

The main results of this section, namely, the modularity (or automorphy) of Calabi–Yau motives, are formulated as follows.

Theorem 9.2. *Let (S, σ) be one of the pairs of the 86 $K3$ surfaces with involution given in* Theorem 4.3. *Let E be an elliptic curve over $\mathbb{Q}$ with involution ι. Let X be a crepant resolution of the quotient threefold $E \times S/\iota \times \sigma$. Then X is a Calabi–Yau threefold with a model defined over $\mathbb{Q}$.*

Moreover, X is modular (automorphic) in the sense of Arthur and Clozel [1]. *That is, the L-series $L(\mathcal{M}_X, \rho, s)$ of the Calabi–Yau motive $\mathcal{M}_X$ is determined by an automorphic representation.*

Specializing further, we have the following result.

Theorem 9.3. *Let (S, σ) be one of the 86 $K3$ surfaces with involution σ listed in* Theorem 4.3. *Let $\tilde{S}$ be the minimal resolution S. Then $\tilde{S}$ is defined over $\mathbb{Q}$, and is of Delsatre type, so it is of CM type. Let E be an elliptic curve E_2 or E_3 with involution ι. Consider the quotient threefold $E \times S/\iota \times \sigma$, and let X be its crepant resolution. Then X is a Calabi–Yau threefold and has a model defined over $\mathbb{Q}$.*

Furthermore, X is of CM type, and it is modular (automorphic) in the sense of Arthur and Clozel, i.e, the Calabi–Yau motive $\mathcal{M}_X$ is automorphic.

Since any elliptic curve E over $\mathbb{Q}$ is modular, without the assumption that E is of CM type, we have the following most general results.

Theorem 9.4. *Let (S, σ) be one of the 86 pairs of $K3$ surfaces with involution σ defined over $\mathbb{Q}$ in* Theorem 4.3. *Let (E, ι) be an elliptic curve defined over $\mathbb{Q}$. Let X be a crepant resolution of the quotient threefold $E \times S/\iota \times \sigma$. Then X has a model defined over $\mathbb{Q}$, and satisfies the following:*

- (a) *Associated to E, there is a cusp form f_E of weight 2 obtained from the 2-dimensional automorphic representation in $\mathfrak{A}(GL_2(\mathbb{A}_{\mathbb{Q}}))$.*
- (b) *Associated to the $K3$ motive $\mathcal{M}_S$ of S, there is the automorphic form obtained from the automorphic induction $I(\mathcal{M}_S)$ in $\mathfrak{A}(GL_{\varphi(d)}(\mathbb{A}_{\mathbb{Q}}))$ for some $d \in \mathbb{N}$.*
- (c) *The Calabi–Yau motive $\mathcal{M}_X$ of the Calabi–Yau threefold X is given by $\mathcal{M}_X = E \otimes \mathcal{M}_S$ and is defined over $\mathbb{Q}$. Associated to $\mathcal{M}_X$, there is the automorphic form $f_E \otimes I(\mathcal{M}_S)$ in $\mathfrak{A}(GL_{2\varphi(d)}(\mathbb{A}_{\mathbb{Q}}))$.*
- (d) *The Calabi–Yau motive $\mathcal{M}_X$ is modular in the sense of Arthur and Clozel.*

Remark 9.1. There are other motives of a Calabi–Yau variety X beside the Calabi–yau motive $\mathcal{M}_X$. The L-series of étale ℓ-adic realizations of these other motives are also modular (automorphic), indeed, they are all induced by the GL_1 automorphic representation of the cyclotomic field $\mathbb{K}$ over $\mathbb{Q}$ in which Jacobi sum Grossencharacters belong. These other motives are more "algebraic" compared with the Calabi–Yau motive $\mathcal{M}_X$.

In the next two sections, we will give proofs to the above theorems. In the next section automorphic inductions.

10. BASE CHANGE AND AUTOMORPHIC INDUCTION

Let F be a number field, and let $\mathbb{A}_F$ be the adele ring of F and $\mathfrak{A}(GL_n(\mathbb{A}_F))$ be the space of automorphic representations of $GL_n(\mathbb{A}_F)$.

Let K be a Galois extension of F with Galois group $G = \mathrm{Gal}(K/F)$. If ρ is an irreducible representation of G, let $L(s, \rho, K/F)$ denote the Artin L-series attached to ρ. The definition of the Artin L-series can be extended to arbitrary representation by multiplicativity of L-series associated to irreducible ones. For each ρ, the Langlands reciprocity conjecture states that there exists an automorphic representation $\pi(\rho) \in \mathfrak{A}(GL_n(\mathbb{A}_F))$ such that

$$L(s, \rho, K/F) = L(s, \pi(\rho)).$$

We now recall a theorem of Arthur and Clozel [1] on base change and automorphic induction for GL_n.

Proposition 10.1 (Arthur–Clozel)**.** *Suppose that K/F is a cyclic extension of prime degree ℓ. Let π and Π denote representations which are induced from cuspidal representation of $GL_n(\mathbb{A}_F)$ and $GL_n(\mathbb{A}_K)$, respectively. Then*

(1) *the base change lift of π, denoted by $B(\pi)$, exists, and it is an automorphic representation in $\mathfrak{A}(GL_n(\mathbb{A}_K))$, and*

(2) *the automorphic induction $I(\Pi)$ of Π, denoted by $I(\Pi)$, exists, and it is an automorphic representation in $\mathfrak{A}(GL_n(\mathbb{A}_F))$.*

Remark 10.1. Arthur–Clozel Theorem can be generalized to any abelian extension K/F.

We can reformulate Arthur–Clozel theorem in terms of L-series. We need to consider Rankin–Selberg L-series of convolution. Let π and π' be two cuspidal, unitary automorphic representations of $GL_n(\mathbb{A}_F)$ and $GL_m(\mathbb{A}_F)$, respectively. Let $\mathfrak{S}$ be a finite set of primes of F such that π and π' are unramified outside $\mathfrak{S}$. Let $L(s, \pi \otimes \pi')$ be the Rankin–Selberg L-series of convolution. Then the result of Arthur and Clozel mentioned above is formulated in terms of L-series as follows:

Lemma 10.2. *Let K/F be cyclic extension of prime degree, and suppose that π and Π are cuspidal unitary automorphic representations of $GL_n(\mathbb{A}_F)$ and $GL_m(\mathbb{A}_K)$, respectively. Then*

$$L(s, B(\pi) \otimes \pi') = L(s, \pi \otimes I(\pi')).$$

Now we describe the L-series of the $K3$ motive $\mathcal{M}_S$ for our $K3$ surfaces S with involution σ, and also the L-series of the Calabi–Yau motive $\mathcal{M}_X$ of a Calabi–Yau threefold X. For this we first need to define Hecke L-series with Jacobi sum Grossencharacters.

We recall now the definition of weighted Jacobi sums and weighted Fermat (diagonal) motives from Gouvêa–Yui [10].

Definition 10.1. (a) Let $\mathbb{K} = \mathbb{Q}(\zeta_d)$ be the d–th cyclotomic field over $\mathbb{Q}$, $\mathcal{O}_{\mathbb{K}}$ the ring of integers of $\mathbb{K}$. Let $\mathfrak{p} \in \mathrm{Spec}(\mathcal{O}_{\mathbb{K}})$. For every $x \in \mathcal{O}_{\mathbb{K}}$ relatively prime to $\mathfrak{p}$, let $\chi_{\mathfrak{p}}(x \bmod \mathfrak{p}) = (\frac{x}{\mathfrak{p}})$ be the d–th power residue symbol on $\mathbb{K}$. If $x \equiv 0 \pmod{\mathfrak{p}}$, we put $\chi_{\mathfrak{p}}(x \bmod \mathfrak{p}) = 0$. Let $(w_0, w_1, w_2, \cdots, w_{n+1})$ be a weight. Define the set

$$\begin{aligned}&\mathfrak{A}_d(w_0, w_1, \cdots, w_{n+1})\\ &:= \left\{\mathbf{a} = (a_0, a_1, \cdots, a_{n+1}) \mid a_i \in (w_i\mathbb{Z}/d\mathbb{Z}), a_i \neq 0, \sum_{i=0}^{n+1} a_i = 0\right\}.\end{aligned}$$

For each $\mathbf{a} \in \mathfrak{A}_d(w_0, w_1, \cdots, w_{n+1})$, the weighted *Jacobi sum* is defined by

$$j_{\mathfrak{p}}(\mathbf{a}) = j_{\mathfrak{p}}(a_0, a_1, \cdots, a_{n+1}) := (-1)^n \sum \chi_{\mathfrak{p}}(v_1)^{a_1}\chi_{\mathfrak{p}}(v_2)^{a_2} \cdots \chi_{\mathfrak{p}}(v_{n+1})^{a_{n+1}}$$

where the sum is taken over $(v_1, v_2, \cdots, v_{n+1}) \in (\mathcal{O}_{\mathbb{K}}/\mathfrak{p})^\times \times \cdots \times (\mathcal{O}_{\mathbb{K}}/\mathfrak{p})^\times$ subject to the linear relation $1 + v_1 + v_2 + \cdots + v_{n+1} \equiv 0 \pmod{\mathfrak{p}}$.

Weighted Jacobi sums are elements of $\mathcal{O}_{\mathbb{K}}$ with complex absolute value equal to $q^{n/2}$ where $q =\mid \mathrm{Norm}\,\mathfrak{p} \mid \equiv 1 \pmod{d}$.

(b) The Galois group $\mathrm{Gal}(\mathbb{K}/\mathbb{Q}) \simeq (\mathbb{Z}/d\mathbb{Z})^\times$ acts on weighted Jacobi sums, by multiplication by $t \in (\mathbb{Z}/d\mathbb{Z})^\times$ on each component of $\mathbf{a}$. Let A denote the $(\mathbb{Z}/d\mathbb{Z})^\times$-orbit of $\mathbf{a}$. For a weighted Jacobi sum $j_{\mathfrak{p}}(\mathbf{a})$, the $(\mathbb{Z}/d\mathbb{Z})^\times$-orbit of $j_{\mathfrak{p}}(\mathbf{a})$ is called the weighted *Fermat (diagonal) motive*, and denoted by $\mathcal{M}_A$.

To each $\mathbf{a} = (a_0, a_1, \cdots, a_{n+1}) \in \mathfrak{A}_d(w_0, w_1, \cdots, w_{n+1})$, define the *length* of $\mathbf{a}$ to be

$$\|\mathbf{a}\| := \left(\frac{1}{d}\sum_{i=0}^{n+1} a_i\right) - 1.$$

Via cohomological realizations of these motives, we can compute the numerical characters of $\mathcal{M}_A$.

- The i-th Betti number is

$$B_i(\mathcal{M}_A) := \dim_{\mathbb{Q}_\ell} H^i(\bar{\mathcal{M}}_A, \mathbb{Q}_\ell) = \begin{cases} \#A & \text{if } i = n \\ 1 & \text{if } i \text{ is even and } A = [0] \\ 0 & \text{otherwise}\end{cases}$$

- The (i, j)-th Hodge nubmer is

$$h^{i,j}(\mathcal{M}_A) := \dim_{\mathbb{C}} H^j(\bar{\mathcal{M}}_A, \Omega^i) = \begin{cases} \#\{\mathbf{a} \in A \mid ||\mathbf{a}|| = i\} & \text{if } i + j = n \\ 1 & \text{if } A = [0] \\ 0 & \text{otherwise}\end{cases}$$

Lemma 10.3. (a) *Let S be a $K3$ surface of degree d in a weighted projective 3-space $\mathbb{P}^3(w_0, w_1, w_2, w_3)$ and suppose that S is of CM type. Then there is the $K3$ motive $\mathcal{M}_S$ associated to the weight $\mathbf{w} = (w_0, w_1, w_2, w_3)$ such that $h^{0,2}(\mathcal{M}_S) = 1$ and $B_2(\mathcal{M}_S) = \varphi(d)$.*

(b) *Let X be a Calabi–Yau threefold of degree d in a weighted projective 4-space $\mathbb{P}^4(w_0, w_1, w_2, w_3, w_4)$, and suppose that X is of CM type. Then there is the Calabi–Yau motive $\mathcal{M}_X$ assocaited to the weight $\mathbf{w} = (w_0, w_1, \cdots, w_4)$ such that $h^{0,3}(\mathcal{M}_X) = 1$ and $B_3(\mathcal{M}_X) = \varphi(d)$.*

In both dimension 2 and 3, all other motives have $h^{0,d}(\mathcal{M}_A) = 0$, that is, they are "algebraic".

Proof. For the weight $\mathbf{w} = (w_0, w_1, \cdots, w_{d+1})$, $\|\mathbf{w}\| = 0$ because of the Calabi–Yau condition: $d = \sum_{i=0}^{d+1} w_i$. Under the assumption that S and X are of CM type, they are realized as finite quotients of some Fermat varieties. So the motive associated to $\mathbf{w}$ has $h^{0,d} = h^{d,0} = 1$. Its dimension is the order of $(\mathbb{Z}/d\mathbb{Z})^\times$, which is $\varphi(d)$ where φ denotes the Euler phi function. All other motives have length ≥ 1. For details on Fermat (diagonal) motives, the reader is referred to Gouvêa–Yui [10]. □

Proposition 10.4. *Let (S, σ) be one of the 86 $K3$ surfaces with involution σ defined in* Theorem 4.3 *by a hypersurface of degree d over $\mathbb{Q}$ in a weighted projective 3-space $\mathbb{P}^3(w_0, w_1, w_2, w_3)$. Then S is of CM type. The L-series of S is determined by the Jacobi sum Grossencharacter of the cyclotomic field $\mathbb{K} := \mathbb{Q}(\zeta_d)$ over $\mathbb{Q}$.*

Let $\mathbf{w} = (w_0, w_1, w_2, w_3)$ be the weight defining S, and let $\mathcal{M}_S$ be the $K3$ motive associated to $\mathbf{w}$. Let $j_{\mathfrak{p}}(\mathbf{w})$ be the Jacobi sum associated to it. Then $j_{\mathfrak{p}}(\mathbf{w})$ is an algebraic integer in $\mathcal{O}_{\mathbb{K}}$ with complex absolute value p. The $K3$ motive $\mathcal{M}_S$ associated to $\mathbf{w}$ is transcendental and corresponds to the single $(\mathbb{Z}/d\mathbb{Z})^\times$-orbit of $j_{\mathfrak{p}}(\mathbf{w})$. Therefore the Galois representation associated to $\mathcal{M}_S$ is induced by a GL_1 automorihic representation of $\mathbb{K} = \mathbb{Q}(\zeta_d)$, and it is irreducible over $\mathbb{Q}$ of dimension $\varphi(d)$. Consequently, the Galois representation of $\mathcal{M}_S$ is the automorphic induction of the GL_1 Grossencharacter representation of $\mathbb{K}$.

In other words, $\mathcal{M}_S$ is modular (automorphic) in the sense of Arthur and Clozel, that is, $L(\mathcal{M}_S, \rho, s)$ is determined by an automorphic representation over $\mathbb{Q}$.

Proof. Our $K3$ surface S is defined by a hypersurface of degree d over $\mathbb{Q}$. S is of CM type. The characteristic polynomial of the Frobenius of the $K3$ motive $\mathcal{M}_S$ has reciprocal root $j_{\mathfrak{p}}(\mathbf{w})$ and their Galois conjugates, that is, the $(\mathbb{Z}/d\mathbb{Z})^\times$-orbit of $j_{\mathfrak{p}}(\mathbf{w})$. We know that $j_{\mathfrak{p}}(\mathbf{w})$ and their Galois conjugates are elements of the cyclotomic field $\mathbb{K} = \mathbb{Q}(\zeta_d)$. The restriction of $\mathrm{Gal}(\bar{\mathbb{Q}}/\mathbb{K})$ is a sum of GL_1-dimensional representations corresponding to Jacobi sum Grossencharacters. So $\mathcal{M}_S$ is G_1-modular over $\mathbb{K}$. Then the Arthur–Clozel automorphic induction process yields the autormorphic representation $I(\mathcal{M}_S)$ in $\mathfrak{A}(GL_{\varphi(d)}(\mathbb{Q}))$, which is irreducible over $\mathbb{Q}$ of dimension $\varphi(d)$. Therefore, $\mathcal{M}_S$ is modular (automorphic) in the sense of Arthur and Clozel. □

11. Proof of the automorphy of our Calabi–Yau threefolds over $\mathbb{Q}$

First we will study in detail the results of Arthur and Clozel [1] about base change and automorphic induction for specific cases which are relevant to our later discussions.

Lemma 11.1. *Let $\mathbb{K} = \mathbb{Q}(\zeta_d)$ be the d-th cyclotomic field over $\mathbb{Q}$. Let ψ be a Jacobi sum Grossencharacter of K. Let $\phi \in \mathfrak{A}(GL_2(\mathbb{A}_{\mathbb{Q}}))$ be an automorphic representation. Then there is the base change representation $B_{K/\mathbb{Q}}(\phi) \in \mathfrak{A}(GL_2(\mathbb{A}_K))$.*

Furthermore, there exists the automorphic induction

$$I(B_{K/\mathbb{Q}}(\phi)) \otimes \psi \in \mathfrak{A}(GL_{2\varphi(d)}(\mathbb{A}_{\mathbb{Q}}).$$

Finally, the Rankin–Selberg L-series is given by

$$L(s, \psi \otimes B_{K/\mathbb{Q}}(\phi)) = L(s, I(B_{K/\mathbb{Q}}(\phi)) \otimes \psi)$$

Proposition 11.2. *Let $d_1, d_2 \in \mathbb{N}$. Let $\mathbb{K}_1 = \mathbb{Q}(\zeta_{d_1})$ and $\mathbb{K}_2 = \mathbb{Q}(\zeta_{d_2})$ be d_1-th and d_2-th cyclotomic fields over $\mathbb{Q}$. Let ψ_1 and ψ_2 be Jacobi sum Grossencharacters of $\mathbb{K}_1$ and $\mathbb{K}_2$, respectively. Then they are respectively automorphic forms in*

$\mathfrak{A}(GL_1(\mathbb{A}_{\mathbb{K}_1}))$ and $\mathfrak{A}(GL_1(\mathbb{A}_{\mathbb{K}_2}))$, respectively. Consider the induced automorphic representation $\psi_1 \otimes \psi_2$.

(a) *If $\mathbb{K}_1 \not\supseteq \mathbb{K}_2$, then $\psi_1 \otimes \psi_2$ corresponds to an automorphic representation in $\mathfrak{A}(GL_1(\mathbb{A}_{\mathbb{K}_1\mathbb{K}_2})$, and*

$$L(s, \psi_1 \otimes \psi_2) = L(s, \psi_1)L(s, \psi_2).$$

(b) *If $\mathbb{K}_1 = \mathbb{K}_2$, then $\psi_1 \otimes \psi_2$ corresponds to the induced automorphic representation $I(\psi_1 \otimes \psi_2)$ in $\mathfrak{A}(GL_2(\mathbb{A}_{\mathbb{K}_1}))$, and*

$$L(s, \psi_1 \otimes \psi_2) = L(s, I(\psi_1 \otimes \psi_2)).$$

(c) *If $\mathbb{K}_1 \supset \mathbb{K}_2$ but $\mathbb{K}_1 \neq \mathbb{K}_2$, then ψ_2 corresponds to a representation of a subgroup, H, of Gal$(\mathbb{K}_1/\mathbb{Q}) := G$, and $\psi_1 \otimes \psi_2$ corresponds to the induced representation in $\mathfrak{A}(GL_1(\mathbb{A}_{\mathbb{K}_1}))$, and*

$$L(s, \psi_1 \otimes \psi_2) = L(s, \psi_1 \otimes \mathit{Ind}_H^G \psi_1).$$

We will recall the modularity of dimension one Calabi–Yau Varieties, namely, elliptic curves defined over $\mathbb{Q}$. We consider elliptic curves E over $\mathbb{Q}$ with involution ι. Then E is modular. More precisely, the Calabi–Yau motive $\mathcal{M}$ is the whole E. So they correspond to a 2-dimensional Galois representation, which is modular. If $E = E_2$ or E_3, then E has CM, and its L-series is determined by Jacobi-sum Grossencharacters of the cylotomic field $\mathbb{Q}(\zeta_{d_1})$ over $\mathbb{Q}$ for some d_1.

For Calabi–Yau varieties of dimension two and their modularity, we confine ourselves to one of the 86 pairs (S, σ) of $K3$ surfaces with involution σ in Theorem 4.3. Then S is of CM type, and the $K3$ motive $\mathcal{M}_S$ is modular (automorphic) in the sense of Arthur and Clozel. (In fact, the whole S is modular (automorphic) by Theorem 4.5.)

Finally we are ready to prove the assertions in Section 9 on the modularity (automorphy) of the Calabi–Yau threefolds of CM type defined over $\mathbb{Q}$. We will present proofs for Theorem 9.3 and Theorem 9.4.

Proof of Theorem 9.3: We can find models (maybe singular) over $\mathbb{Q}$ which are hypersurface defining equations for our Calabi–Yau threefolds. That X is of CM type is proved in Theorem 7.2 for the 45 Calabi–Yau threefolds over $\mathbb{Q}$ where the $K3$ components S are of Borcea type. For the remaining 41 cases, this is proved in Theorem 7.3. When X is of CM type, the Galois representation of X is induced from the one-dimensional representations associated to the components, and the automorphic induction process yields irreducible automorphic representation over $\mathbb{Q}$. Therefore X is modular (automporpphic). □

Proof of Theorem 9.4:

(a) This follows from the modularity theorem for elliptic curves defined over $\mathbb{Q}$. Associated to E, there is a cusp form f_E of weight 2 obtained from the 2-dimensional automorphic representation in $\mathfrak{A}(GL_2(\mathbb{A}_{\mathbb{Q}}))$.

(b) Theorem 4.5 is valid for the $K3$ motive $\mathcal{M}_S$. Associated to the $K3$ motive $\mathcal{M}_S$ of S, there is the automorphic form obtained from the automorphic induction $I(\mathcal{M}_S)$ in $\mathfrak{A}(GL_{\varphi(d)}(\mathbb{A}_{\mathbb{Q}}))$ for some $d \in \mathbb{N}$.

(c) The Calabi–Yau motive $\mathcal{M}_X$ of the Calabi–Yau threefold X is given by $\mathcal{M}_X = E \otimes \mathcal{M}_S$ and is defined over $\mathbb{Q}$. Associated to $\mathcal{M}_X$, there is the automorphic form $f_E \otimes I(\mathcal{M}_S)$ in $\mathfrak{A}(GL_{2\varphi(d)}(\mathbb{A}_{\mathbb{Q})}))$. This follows from Lemma 11.1.

(d) The Calabi–Yau motive $\mathcal{M}_X$ is modular (automorphic) in the sense of Arthur and Clozel. This is nothing but the automorphic induction.

□

Remark 11.1. The automorphy of other motives can also be established in the same way as for the Calabi–Yau motive, but indeed much simpler. In fact, the L-series can be expressed in terms of Jacobi sum Grossencharacters.

Problem 2. How to give explicit descriptions for the automorphic forms associated to the Calabi–Yau motive $\mathcal{M}_X$ remains an open question.

12. The automorphy of mirror Calabi–Yau threefolds over $\mathbb{Q}$

Mirror symmetry for $K3$ surfaces is given by Arnold's strange duality. It is due to Arnold–Dolgachev–Nikulin and is formulated for lattice polarized $K3$ surfaces as follows: A pair of lattice polarized $K3$ surfaces $(S, S^\vee)$ is said to be a mirror pair if

$$\mathrm{Pic}\,(S)^{\perp}_{H^2(S,\mathbb{Z})} = U \oplus \mathrm{Pic}\,(S^\vee)$$

as lattices. In terms of the Picard numbers,

$$22 - \rho(S) = 2 + \rho(S^\vee) \quad \Leftrightarrow \quad \rho(S^\vee) = 20 - \rho(S).$$

(See Dolgachev [5] for details.)

Now we consider mirror symmetry for pairs (S, σ) of $K3$ surfaces with involution σ classified by Nikulin in terms of triplets (r, a, δ). Let (S, σ) be a pair of $K3$ surfaces with involution σ corresponding to a triplet (r, a, δ). Then the mirror pair $(S^\vee, \sigma)$ corresponds to the triplet $(20 - r, a, \delta)$.

For the Nikulin pyramid given in Section 2, the mirror is placed at the vertical line $r = 10$, corresponding to the symmetry $(r, a, \delta) \leftrightarrow (20 - r, a, \delta)$. It should be remarked that mirrors do not exist for the points located at the utmost right outerlayer of the pyramid, that is, (r, a, δ) is one of the following triplets $(20, 2, 1)$, $(19, 3, 1)$, $(18, 4, 1)$, $(18, 4, 0)$, $(17, 5, 1)$, $(16, 6, 1)$, $(15, 7, 1)$, $(14, 8, 1)$, $(13, 9, 1)$ and $(12, 10, 1)$.

Now we consider our Calabi–Yau threefolds obtained as crepant resolutions of quotient threefolds $E \times S/\iota \times \sigma$. Mirror symmetry for these Calabi–Yau families has been discussed by Voisin [20] and also by Borcea [4].

Theorem 12.1. *Given a Calabi–Yau family $X = X(r, a, \delta) = E \times \widetilde{S/\iota} \times \sigma$, there is a mirror Calabi–Yau family $X^\vee = X(20 - r, a, \delta) = E \times \widetilde{S^\vee/\iota} \times \sigma$ such that*

$$e(E^\vee) = -e(X)$$

Mirror symmetry for Calabi–Yau threefolds X is purely determined by mirror symmetry for the $K3$ components S.

Borcea's formulation of mirror symmetry is:

$$h^{1,1}(X^\vee) = 5 + 3(20 - r) - 2a = 65 - 3r - 2a = h^{2,1}(X),$$

$$h^{2,1}(X^\vee) = 65 - 3(20 - r) - 2a = 5 + 3r - 2a = h^{1,1}(X)$$

and

$$e(X^\vee) = -12(r - 10) = -e(X)$$

That is, mirror symmetry interchanges r by $20 - r$.

Voisin's formulation of mirror symmetry is given as follows: Recall that the fixed part S^σ of S under σ ia a disjoint union of a genus-g curve and k rational curves on S. Put

$$N := 1 + k = \text{the number of components of } S^\sigma$$
$$N' := \text{the sum of genera of components of } S^\sigma$$

Then

$$h^{1,1}(X) = 11 + 5N - N',$$
$$h^{2,1}(X) = 11 + 5N' - N$$

and

$$e(X) = 12(N - N').$$

Mirror symmetry interchanges N and N'.

$$h^{1,1}(X^\vee) = 11 + 5N' - N$$
$$h^{2,1}(X^\vee) = 11 + 5N - N'$$

and

$$e(X^\vee) = 12(N' - N) = -e(X)$$

Remark 12.1. For the Calabi–Yau threefolds corresponding to the 10 triplets (r, a, δ) located at the utmost right outerlayer of the Nikulin's pyramid (called the "pale region" by Borcea), mirror partners do not exist. This was recently pointed out by Rohde [19], and studied by Garbagnati–van Geemen [8]. A reason for this is the non-existence of boundary points in the complex structure moduli space of the Calabi–Yau threefold where the variation of Hodge structures on H^3 has maximal unipotent monodromy.

Now we consider the 95 families of $K3$ surfaces in the list of Reid and Yonemura. Belcastro [2] determined the Picard lattices for these 95 $K3$ surfaces, and showed that these 95 families are not closed under mymmetry.

We can fish out those $K3$ surfaces with involution σ which are closed under mirror symmetry.

Lemma 12.2 (Belcastro [2])**.** *The* 95 *families of* $K3$ *surfaces of Reid and Yonemura are not closed under mirror symmetry. Among them, the* 57 *families have mirror partners within the list.*

Lemma 12.3. *All* 57 *families of* $K3$ *surfaces* S *have non-symplectic involutions* σ *acting as* -1 *on* $H^2(S)$*, and their mirror partners* $S^\vee$ *also non-symplective involutions* σ *with the same property.*

Proof. We tabulate the 57 families of $K3$ surfaces with involutions, and their mirror partners in Table 10 and Table 11. □

Theorem 12.4. *Let* (S, σ) *be one of the* 57 $K3$ *surfaces with involution* σ*, which are closed under mirror symmetry. Let* X *be a Calabi–Yau threefold corresponding to a triplet* (r, a, δ)*. Then a mirror Calabi–Yau threefold* $X^\vee$ *exists and corresponds to the triplet* $(20 - r, a, \delta)$*. Then* $X^\vee$ *is obtained as a crepant resolution of the quotient* $E \times S^\vee/\iota \times \sigma$*.* $X^\vee$ *satisfies the following:*

- (a) $X^\vee$ *has a model defined over* $\mathbb{Q}$
- (b) $X^\vee$ *is of CM type if and only if* E *is of CM type.*
- (c) *If* $X^\vee$ *is of CM type, then* X *is modular (automorphic) in the sense of Arthur and Clozel.*

Proof. Repeat the same proof as for Theorem 9.4 to $X^\vee$. $\square$

Example 12.5. Let $E = E_2$ be an elliptic curve with involution ι and let $S = \#14$ in Yonemura $=\#26$ in Borcea be a $K3$ surface given by

$$S : x_0^2 = x_1^3 + x_2^7 + x_3^{42} \subset \mathbb{P}^3(21, 14, 6, 1)$$

of degree 42 and involution $\sigma(x_0) = -x_0$. S corresponds to the triplet $(10, 0, 0)$. A Calabi–Yau threefold $X = E_2 \widetilde{\times S/\iota} \times \sigma$ has a birational model defined over $\mathbb{Q}$

$$X : z_0^4 + z_1^4 = z_2^3 + z_3^7 + z_4^{42} \subset \mathbb{P}^4(21, 21, 28, 12, 2)$$

of degree 84. The Hodge numbers and the Euler characteristic are

$$h^{1,1}(X) = 35,\ h^{2,1}(X) = 35,\ e(X) = 0$$

so that X is its own mirror.

The Calabi–Yau motive $\mathcal{M}_X$ has dimension $\varphi(84) = 24$ and the Jacobi sum Grossencharacter of $K = \mathbb{Q}(\zeta_{84})$ gives GL_1-representations and its automorphic induction gives rise to the GL_{24} irreducible representation for $\mathcal{M}_X$ over $\mathbb{Q}$, and hence it is modular (automorphic).

Example 12.6. Let $E = E_2$ be an elliptic curve with involution ι and let $S = \#40$ in Yonemura $=\#5$ in Borcea be a $K3$ surface given by

$$S : x_0^2 = x_1^3 x_2 + x_1^3 x_3^2 + x_2^7 - x_3^{14} \subset \mathbb{P}^2(7, 4, 2, 1)$$

of degree 14 with involution $\sigma(x_0) = -x_0$. S corresponds to the triplet $(7, 3, 0)$. By Theorem 4.3, we may remove the monomial $x_1^3 x_2^2$ from the defining equation.

A Calabi–Yau threefold X has a birational model defined over $\mathbb{Q}$:

$$X : z_0^4 + z_1^4 = z_2^3 z_3 + z_3^7 - z_4^{14} \subset \mathbb{P}^4(7, 7, 8, 4, 2)$$

of degree 28. The Hodge numbers and the Euler characteristic are

$$h^{1,1}(X) = 20,\ h^{2,1}(X) = 38,\ e(X) = -36.$$

The Calabi–Yau motive $\mathcal{M}_X$ has dimension $\varphi(84) = 24$ and the Jacobi Grossencharacter of $\mathbb{K} = \mathbb{Q}(\zeta_{84})$ gives GL_1 representations and its automorphic induction gives rise to GL_{24} irreducible representation for $\mathcal{M}_X$ over $\mathbb{Q}$. Hence $\mathcal{M}_X$ is modular (automorphic).

A candidate for mirror partner for X is $E_2 \widetilde{\times S^\vee/\iota} \times \sigma$ where $S^\vee$ may be taken to be $\#47$ in Yonemura $=\#24$ in Borcea. $S^\vee$ is a $K3$ surface defined by

$$S^\vee : x_0^2 = x_1^3 + x_1 x_2^7 + x_2^9 x_3^2 + x_3^{14} \subset \mathbb{P}^3(21, 14, 4, 3)$$

of degree 42. $S^\vee$ corresponds to the triplet $(13, 3, 0)$. By Theorem 4.3, we may remove the monomial $x_2^9 x_3^2$ from the defining equation. A mirror Calabi–Yau threefold $X^\vee$ has a birational model defined over $\mathbb{Q}$:

$$X^\vee : z_0^4 + z_1^4 = z_2^3 + z_2 z_3^7 + z_4^{14} \subset \mathbb{P}^4(21, 21, 28, 8, 6)$$

of degree 84. The Hodge numbers and the Euler characteristic of $X^\vee$ are

$$h^{1,1}(X^\vee) = 38,\ h^{2,1}(X^\vee) = 20,\ e(X^\vee) = 36.$$

The Calabi–Yau motive $\mathcal{M}_{X^\vee}$ has dimension $\varphi(28) = 12$. Again, the Jacobi sum Grossencharacter of $\mathbb{K} = \mathbb{Q}(\zeta_{28})$ gives GL_1 representations and its automorphic induction yields GL_{12} irreducible representation for $\mathcal{M}_{X^\vee}$ over $\mathbb{Q}$. So $\mathcal{M}_{X^\vee}$ is modular (automorphic).

For these two examples, it happens that $L(\mathcal{M}_X, \rho, s) = L(\mathcal{M}_{X^\vee}, \rho^\vee, s)$. That is the L-series of the Calabi–Yau motives of X and $X^\vee$ coinside.

Example 12.7. Let $E = E_3$ be an elliptic curve with involution ι and let $S = \#80$ in Yonemura $= \#40$ in Borcea be a $K3$ surface given by

$$S : x_0^2 = x_1^3x_2 + x_2^8x_3 + x_3^{11} \subset \mathbb{P}^3(22, 13, 5, 4)$$

of degree 44 with involution $\sigma(x_0) = -x_0$. S corresponds to the triplet $(18, 0, 0)$. S is already of CM type.

A Calabi–Yau threefold X has a birational model defined over $\mathbb{Q}$:

$$X : z_0^3 + z_1^6 = z_2^3z_3 + z_3^8z_4 + z_4^{11} \subset \mathbb{P}^4(44, 22, 39, 15, 12)$$

of degree 132. The Hodge numbers and the Euler characteristic are

$$h^{1,1}(X) = 59,\ h^{2,1}(X) = 11,\ e(X) = 96.$$

The Calabi–Yau motive $\mathcal{M}_X$ has dimension $\varphi(12) = 4$. The Jacobi sum Grossencharacter of $\mathbb{K} = \mathbb{Q}(\zeta_{12})$ gives GL_1 representations and its automorphic induction yields GL_4 irreducible representation for $\mathcal{M}_X$ over $\mathbb{Q}$. So $\mathcal{M}_X$ is modular (automorphic).

A candidate for mirror partner for X is the product $E \times S^\vee$ where $S\vee$ is $\#10$ in Yonemura $= \#42$ in Borcea, which is given by

$$S^\vee : x_0^2 = x_1^3 + x_2^{12} + x_3^{12} \subset \mathbb{P}^3(6, 4, 1, 1)$$

of degree 12. However, since w_0 is divisible by 6, we do not know how to construct a hypersurface defining equation for a mirror Calabi–Yau threefold $X^\vee$.

Problem 3. Let $(X, X^\vee)$ be a mirror pair of Calabi–Yau threefolds of Theorem 12.1. As the first two examples indicates, the L-seires of the Calabi–Yau motive $\mathcal{M}_X$ remain invariant under mirror symmetry, so that $L(\mathcal{M}_X, \rho, s) = L(\mathcal{M}_{X^\vee}, \rho^\vee, s)$. A natural question is: Does the Calabi–Yau motive $\mathcal{M}_X$ remain invariant under mirror symmetry?

13. TABLES

Some clarifications might be in order how to read the tables.

- In the Table 1, 2 and 3, we use two numbering systems, one from Borcea $B\#$ and the other from Yonemura $Y\#$. We matched up the numbers in two lists.
- We use two sets of notations for variables, one is $x_0, x_1, \cdots, x_n$, and the other is $x, y, z, w, \cdots$. In relevant tables, we indicated identification of these two sets of variables.
- The equations in Tables 1, 2 and 3 are taken from Borcea's paper [4]. "Terms removed" indicates the terms we may remove to specialize the equations (or deformation) into Fermat or Delsarte type.
- The equations in Table 4 are taken from Yonemura's paper [21]. "Terms removed" indicates the terms we may remove to specialize the equations into Delsarte type. In Case #1, there is no choice of equations of the form $x_0^2 = f(x_1, x_2, x_3)$ or $x_0^2x_i = f(x_1, x_2, x_3)$.
- In Table 5, the equations are taken from Yonemura's paper [21]. In order to make the equations into Delsarte type, we slightly generalize the original equations and then remove some terms, if necessary. In other words, we first

add a few terms to the original equation and then remove several terms to make the equation into a form of Delsarte type. This procedure is indicated as "terms changed/removed."

For instance, in the case #17, we first add a term x^2y and then remove x^3, xw^5 and yw^5. In effect, this procedure interchanges x^3 with x^2y, and remove xw^5 and yw^5. It results in a new equation $x^2y + y^3 + z^5 + zw^6$.

- In Table 6, the equations are taken from Yonemura's paper [21]. For the $K3$ surfaces on this table, there is no way to define the involution $\sigma(x) = -x$ by using equations of Delsarte type. We therefore define an involution on some other variable. This alternative involution is indicated in the column "involution." Nikulin's invariants r and a for such (S, σ) are calculated in Table 9.
- The $K3$ surfaces in Table 7 are taken from Yonemura's paper [21]. They have involution $\sigma(x) = -x$, but no matter what terms we add or remove from the equations, we cannot transform them into quasi-smooth equations of Delsarte type (cf. Remark 4.4).
- The $K3$ surfaces in Table 8 are also taken from Yonemura's paper [21]. For them, we do not know how to define a non-symplectic involution on the $K3$ surfaces by preserving their quasi-smoothness (even if we allow more than four monomials in the equations).
- Table 9 lists $K3$ surfaces with involution on x_1, x_2 or x_3 (i.e., not the variable x_0 of highest weight). The variable we choose is indicated under the column $\sigma(x_i) = -x_i$. For some $K3$ surfaces, we consider two involutions.

Acknowledgments

Y. Goto is supported in part by JSPS grant from Japan (category C, 21540003 and 24540004). R. Livné is supprted in part by a grant from Israel. N. Yui is supported in part by NSERC Discovery grant from Canada. The paper is completed while N. Yui was a visiting professor at Kavli IPMU, and at Tsuda College in Japan. She thanks the hospitality of the two institutions. Y. Goto prepared the manuscript while he visited the Fields Institute and Queen's University in Canada, Tsuda College in Japan and the National Center for Theoretical Sciences of Taiwan. He thanks the hospitality of these institutions.

References

[1] Arthur, J., and Clozel, L., *Simple algebras, base change and the advanced theory of the trace formula* Annals of Math. Stud. **120**, Princeton University Press 1990.

[2] Belcastro, S.-M., *Picard lattices of families of K3 surfaces*, Commun. in Algebra **30** (2002), pp. 61–82. arXiv:math.AG/9809008

[3] Borcea, C., *Calabi–Yau manifolds and complex multiplciation*, in *Essays on mirror manifolds*, International Press, Hong Kong 1992, pp. 489–502.

[4] Borcea, C., *K3 surfaces with involution and mirror pairs of Calabi–yau manifolds*, in *Mirror Symmetry*, AMS/IP Stud. Adv. Math. **1**, American Mathematical Society, Providence, RI 1997, pp. 717–743.

[5] Dolgachev, I., *Mirror symmetry for lattice polarized K3 surfaces*, J. Math. Sci. **81** (1996), pp. 2599–2630.

Table 1. $K3$ weights in Borcea's list with odd w_0

$Y\#$	$B\#$	(w_0,w_1,w_2,w_3)	$f(x_1,x_2,x_3)=f(y,z,w)$	r	a	terms removed from equations of [B]
5	1	$(3,1,1,1)$	$y^6+z^6+w^6$	1	1	
6	2	$(5,2,2,1)$	$y^5+z^5+w^{10}$	6	4	
42	3	$(5,3,1,1)$	$y^3z+z^{10}+w^{10}$	3	1	y^3w
32	4	$(7,3,2,2)$	$y^4z+z^7+w^7$	10	6	y^4w
40	5	$(7,4,2,1)$	$y^3z+z^7+w^{14}$	7	3	y^3w^2
33	6	$(9,4,3,2)$	$y^4w+z^6+w^9$	10	6	y^3z^2
39	7	$(9,5,3,1)$	$y^3z+z^6+w^{18}$	7	3	y^3w^3
12	8	$(9,6,2,1)$	$y^3+z^9+w^{18}$	6	2	
75	9	$(11,5,4,2)$	$y^4w+z^5w+w^{11}$	13	5	y^2z^3
78	10	$(11,6,4,1)$	$y^3z+yz^4+w^{22}$	10	2	y^3w^4, z^5w^2
82	11	$(11,7,3,1)$	$y^3w+yz^5+w^{22}$	9	1	z^7w
76	12	$(13,6,5,2)$	$y^4w+yz^4+w^{13}$	14	4	z^4w^3
77	13	$(13,7,5,1)$	$y^3z+z^5w+w^{26}$	11	1	y^3w^5
81	14	$(13,8,3,2)$	$y^3w+yz^6+w^{13}$	13	3	z^8w
29	15	$(15,6,5,4)$	$y^5+z^6+yw^6$	12	6	z^2w^5
34	16	$(15,7,6,2)$	$y^4w+z^5+w^{15}$	14	4	
38	17	$(15,8,6,1)$	$y^3z+z^5+w^{30}$	11	1	y^3w^6
11	18	$(15,10,3,2)$	$y^3+z^{10}+w^{15}$	10	4	
50	19	$(15,10,4,1)$	$y^3+yz^5+w^{30}$	9	1	z^7w^2
90	20	$(17,7,6,4)$	$y^4z+y^2w^5+z^5w+zw^7$	17	3	no Delsarte form
93	21	$(17,10,4,3)$	$y^3z+yz^6+yw^8+z^7w^2+zw^{10}$	16	2	no Delsarte form
91	22	$(19,8,6,5)$	$y^4z+yz^5+yw^6+z^3w^4$	18	2	no Delsarte form
92	23	$(19,11,5,3)$	$y^3z+yw^9+z^7w$	17	1	zw^{11}
47	24	$(21,14,4,3)$	$y^3+yz^7+w^{14}$	13	3	z^9w^2
49	25	$(21,14,5,2)$	$y^3+z^8w+w^{21}$	14	2	
14	26	$(21,14,6,1)$	$y^3+w^7+w^{42}$	10	0	
73	27	$(25,10,8,7)$	$y^5+yz^5+zw^6$	19	1	
83	28	$(27,18,5,4)$	$y^3+yw^9+z^{10}w$	17	1	z^2w^{11}
46	29	$(33,22,6,5)$	$y^3+z^{11}+zw^{12}$	18	0	

[6] Dolgachev, I., and Kondo, S., *The rationality of the moduli spaces of Coble surfaces and of nodal Enriques surfaces*, arXiv:1201.6093.

[7] Fletcher, A.R., *Working with weighted complete intersection.*

[8] Garbagnati, A., and van Geemen, B., *Examples of Calabi–Yau threefolds parametrized by Shimura varieties*, Rend. Sem. Mat. Univ. Pol. Torino **68** (2010), no. 3, pp. 271–287.

[9] Goto, Y., Kloosterman, R., and Yui, N., *Zeta-functions of certain $K3$-fibered Calabi–Yau threefolds*, International J. Math, **22**, No. 1 (2011), pp. 67–129.

[10] Gouvêa, F., and Yui, N., *Arithmetic of diagonal hypersurfaces over finite fields*, London Math. Soc. Lecture Notes 209, xii+169 pages, Cambridge University Press, 1995.

[11] Kondo, S., *The rationality of the moduli spaces of Enriques surfaces*, Compositio Math. **91** (1994), pp. 159–173.

[12] Livné, R., and Yui, N., *The modularity of certain non-rigid Calabi–Yau threefolds*, J. Math. Kyoto Univ. **45**, No. 4 (2005), pp. 645–665.

Table 2. $K3$ weights in Borcea's list with even w_0

$Y\#$	$B\#$	(w_0, w_1, w_2, w_3)	$f(x_1, x_2, x_3) = f(y, z, w)$	r	a	terms removed from equations of [B]
7	30	$(4, 2, 1, 1)$	$y^4 + z^8 + w^8$	2	2	
37	31	$(8, 4, 3, 1)$	$y^4 + yz^4 + w^{16}$	6	4	z^5w
44	32	$(8, 5, 2, 1)$	$y^3w + z^8 + w^{16}$	6	2	y^2z^3
36	33	$(10, 5, 3, 2)$	$y^4 + yz^5 + w^{10}$	8	6	z^6w
9	34	$(10, 5, 4, 1)$	$y^4 + z^5 + w^{20}$	6	4	
35	35	$(14, 7, 4, 3)$	$y^4 + z^7 + yw^7$	10	6	zw^8
45	36	$(14, 9, 4, 1)$	$y^3w + z^7 + w^{28}$	10	0	
74	37	$(16, 7, 5, 4)$	$y^4w + yz^5 + w^8$	14	4	z^4w^3
79	38	$(16, 9, 5, 2)$	$y^3z + z^6w + w^{16}$	14	2	y^2w^7
30	39	$(20, 8, 7, 5)$	$y^5 + z^5w + w^8$	14	4	
80	40	$(22, 13, 5, 4)$	$y^3z + z^8w + w^{11}$	18	0	

Table 3. $K3$ weights in Borcea's list with w_0 divisible by 6

$Y\#$	$B\#$	(w_0, w_1, w_2, w_3)	$f(x_1, x_2, x_3) = f(y, z, w)$	r	a	terms removed from equations of [B]
8	41	$(6, 3, 2, 1)$	$y^4 + z^6 + w^{12}$	4	4	
10	42	$(6, 4, 1, 1)$	$y^3 + z^{12} + w^{12}$	2	0	
31	43	$(12, 5, 4, 3)$	$y^4z + z^6 + w^8$	14	2	y^3w^3
41	44	$(12, 7, 3, 2)$	$y^3z + z^8 + w^{12}$	10	4	y^2w^5
13	45	$(12, 8, 3, 1)$	$y^3 + z^8 + w^{24}$	6	2	
43	46	$(18, 11, 4, 3)$	$y^3w + z^9 + w^{12}$	14	2	
51	47	$(18, 12, 5, 1)$	$y^3 + z^7w + w^{36}$	10	0	
48	48	$(24, 16, 5, 3)$	$y^3 + z^9w + w^{16}$	14	2	

[13] Livné, R., Schütt, M., and Yui, N., *The modularity of $K3$ surfaces with non-symplectic group actions*, Math. Annalen **348** (2010), pp. 333–355.

[14] Ma, S., *Rationality of the moduli spaces of 2-elementary $K3$ surfaces, I*, arXiv:1110.5110. 2011.

[15] Ma, S., *Rationality of the moduli spaces of 2-elementary $K3$ surfaces, II*, arXiv;1205.4784, 2012.

[16] Nikulin, V., *Integral symmetric bilinear forms and some of their geometric applications*, Izv. Akad. Nauk SSSR Ser. Mat. bf 43 (1979) = Math. USSR-Izv. **14** (1980), pp. 103–167.

[17] Nikulin, V., *Discrete reflection groups in Lobachevsky spaces and algebraic surfaces*, Proc. ICM Berkeley, California, 1986, pp. 654–671.

[18] Reid, M., *Canonical 3-folds*, Proc. Alg. Geom. Anger 1979. Sijthoff and Nordhoff, pp. 273–310.

[19] Rohde, C., *Cyclic coverings, Calabi–yau manifolds and complex multiplcation*, Lecture Notes in Math. **1975**, Springer 2009.

[20] Voisin, C., *Mirrors et involutions sur les surfaces $K3$*, In *Journées de Géométrie Algeébrique d'Orsay*, Astérisque **218** (1993), pp. 273–323.

[21] Yonemura, H., *Hypersurface simple $K3$ singularities*, Tôhoku J. Math. **42** (1990), pp. 351–380.

Table 4. Delsarte-type $K3$ surfaces with involutions $\sigma(x) = -x$, NOT in Borcea's list, after removal of several terms

$Y\#$	(w_0, w_1, w_2, w_3)	$F(x_0, x_1, x_2, x_3) = F(x, y, z, w)$	r	a	terms removed from equations of [Y]
1	$(1, 1, 1, 1)$	$x^4 + y^4 + z^4 + w^4$	8	8	
19	$(3, 2, 2, 1)$	$x^2y + y^4 + z^4 + w^8$	10	6	x^2w^2 and x^2z
20	$(9, 8, 6, 1)$	$x^2z + y^3 + z^4 + w^{24}$	10	6	x^2w^6
21	$(2, 1, 1, 1)$	$x^2y + y^5 + z^5 + w^5$	6	4	x^2z, x^2w
22	$(6, 5, 3, 1)$	$x^2z + y^3 + z^5 + w^{15}$	10	4	x^2w^3
23	$(5, 3, 2, 2)$	$x^2z + y^4 + z^6 + w^6$	12	6	x^2w
24	$(5, 4, 2, 1)$	$x^2z + y^3 + z^6 + w^{12}$	10	4	x^2w^2
25	$(4, 3, 1, 1)$	$x^2z + y^3 + z^9 + w^9$	6	2	x^2w
26	$(9, 5, 4, 2)$	$x^2w + y^4 + z^5 + w^{10}$	14	4	
27	$(11, 8, 3, 2)$	$x^2w + y^3 + z^8 + w^{12}$	14	2	
28	$(10, 7, 3, 1)$	$x^2w + y^3 + z^7 + w^{21}$	11	1	
55	$(7, 6, 5, 2)$	$x^2y + y^3w + z^4 + w^{10}$	14	4	x^2w^3
56	$(11, 8, 6, 5)$	$x^2y + y^3z + z^5 + w^6$	19	1	
57	$(9, 6, 5, 4)$	$x^2y + y^4 + z^4w + w^6$	18	2	xz^3
58	$(6, 5, 4, 1)$	$x^2z + y^3w + z^4 + w^{16}$	14	2	x^2w^4, xy^2
59	$(8, 7, 5, 1)$	$x^2z + y^3 + z^4w + w^{21}$	14	2	x^2w^5
60	$(7, 6, 4, 1)$	$x^2z + y^3 + yz^3 + w^{18}$	13	3	x^2w^4, z^4w^2
61	$(11, 7, 6, 4)$	$x^2z + y^4 + z^4w + w^7$	18	2	
62	$(8, 5, 4, 3)$	$x^2z + y^4 + yw^5 + z^5$	14	4	xw^4, z^2w^4
63	$(4, 3, 2, 1)$	$x^2z + y^3w + z^5 + w^{10}$	10	4	x^2w^2, xy^2, y^2z^2
64	$(10, 7, 4, 3)$	$x^2z + y^3w + z^6 + w^8$	18	0	xy^2
65	$(14, 11, 5, 3)$	$x^2z + y^3 + z^6w + w^{11}$	18	0	
66	$(3, 2, 1, 1)$	$x^2z + y^3w + z^7 + w^7$	7	3	x^2w, xy^2, y^3z
67	$(9, 7, 3, 2)$	$x^2z + y^3 + yw^7 + z^7$	13	3	xw^6, zw^9
68	$(13, 10, 4, 3)$	$x^2z + y^3 + yz^5 + w^{10}$	17	1	z^6w^2
69	$(7, 4, 3, 2)$	$x^2w + y^4 + yz^4 + w^8$	14	4	xz^3, z^4w^2
70	$(8, 5, 3, 2)$	$x^2w + y^3z + z^6 + w^9$	14	2	xy^2, y^2w^4
71	$(7, 4, 3, 1)$	$x^2w + y^3z + z^5 + w^{15}$	11	1	xy^2, y^3w^3
72	$(7, 5, 2, 1)$	$x^2w + y^3 + yz^5 + w^{15}$	9	1	xz^4, z^7w
86	$(9, 7, 5, 4)$	$x^2y + y^3w + z^5 + zw^5$	19	1	xw^4
87	$(5, 4, 3, 1)$	$x^2z + y^3w + yz^3 + w^{13}$	13	3	x^2w^3, xy^2, z^4w
88	$(11, 9, 5, 2)$	$x^2z + y^3 + yw^9 + z^5w$	17	1	xw^8, zw^{11}
89	$(5, 3, 2, 1)$	$x^2w + y^3z + yz^4 + w^{11}$	10	2	xy^2, xz^3, y^3w^2, z^5w

[22] Zarhin, Yu., *Hodge groups of $K3$ surfaces*, J. Reine Angew. Math. **341** (1983), pp. 193–220.

Table 5. $K3$ surfaces with involution $\sigma(x) = -x$, NOT in Borcea's list, after change and/or removal of several terms

$Y\#$	(w_0, w_1, w_2, w_3)	$F(x_0, x_1, x_2, x_3) = F(x, y, z, w)$	r	a	terms changed/removed from equations of [Y]
3	$(2, 2, 1, 1)$	$x^2y + y^3 + z^6 + w^6$	7	7	$x^3 \to x^2y$
4	$(4, 4, 3, 1)$	$x^2y + y^3 + z^4 + w^{12}$	7	7	$x^3 \to x^2y$
17	$(5, 5, 3, 2)$	$x^2y + y^3 + z^5 + zw^6$	12	6	$xw^5, yw^5, x^3 \to x^2y$
18	$(3, 3, 2, 1)$	$x^2y + y^3 + z^4w + w^9$	10	6	$xz^3, yz^3, x^3 \to x^2y$

Table 6. $K3$ surfaces with a different kind of involution

$Y\#$	(w_0, w_1, w_2, w_3)	$F(x_0, x_1, x_2, x_3) = F(x, y, z, w)$	terms removed	involution
2	$(4, 3, 3, 2)$	$x^3 + y^4 + z^4 + w^6$	none	$y \to -y$
16	$(8, 7, 6, 3)$	$x^3 + y^3w + z^4 + w^8$	none	$z \to -z$
52	$(12, 9, 8, 7)$	$x^3 + y^4 + xz^3 + zw^4$	none	$y \to -y$
84	$(9, 7, 6, 5)$	$x^3 + xz^3 + y^3z + yw^4$	z^2w^3	$w \to -w$

Table 7. $K3$ surfaces with involution $\sigma(x) = -x$, but not realized as quasi-smooth hypersurfaces in 4 monomials

$Y\#$	(w_0, w_1, w_2, w_3)	$F(x_0, x_1, x_2, x_3) = F(x, y, z, w)$	r	a
85	$(5, 4, 3, 2)$	$x^2y + x^2w^2 + y^3w + y^2z^2 + yw^5 + z^4w + w^7$	15	5
90	$(17, 7, 6, 4)$	$x^2 + y^4z + y^2w^5 + z^5w + zw^7$	17	3
91	$(19, 8, 6, 5)$	$x^2 + y^4z + yz^5 + yw^6 + z^3w^4$	18	2
93	$(17, 10, 4, 3)$	$x^2 + y^3z + yz^6 + yw^8 + z^7w^2 + zw^{10}$	16	2
94	$(7, 5, 4, 3)$	$x^2y + y^3z + y^2w^3 + z^4w + zw^5$	18	2
95	$(7, 5, 3, 2)$	$x^2z + y^3w + yz^4 + yw^6 + z^5w + zw^7$	16	2

Table 8. $K3$ weights with no obvious involution

$Y\#$	(w_0, w_1, w_2, w_3)	$F(x_0, x_1, x_2, x_3) = F(x, y, z, w)$
15	$(5, 4, 3, 3)$	$x^3 + y^3z + y^3w + z^5 + w^5$
53	$(6, 5, 4, 3)$	$x^3 + y^3w + y^2z^2 + xz^3 + z^3w^2 + w^6$
54	$(7, 6, 5, 3)$	$x^3 + y^3w + yz^3 + z^3w^2 + w^7$

Department of Mathematics, Hokkaido University of Education, 1-2 Hachiman-cho, Hakodate 040-8567 Japan
E-mail address: `goto.yasuhiro@h.hokkyodai.ac.jp`

Institute of Mathematics, Hebrew University of Jerusalem Givat-Ram, Jerusalem 919004, Israel
E-mail address: `rlivne@math.huji.ac.il`

Department of Mathematics and Statistics, Queen's University, Kingston, Ontario Canada K7L 3N6
E-mail address: `yui@mast.queensu.ca`

Table 9. Nikulin's invariant associated with other types of involutions

$Y\#$	(w_0,w_1,w_2,w_3)	$F(x_0,x_1,x_2,x_3)=F(x,y,z,w)$	$\sigma(x_i)=-x_i$	r	a
2	$(4,3,3,2)$	$x^3+y^4+z^4+w^6$	y	10	8
	$(4,3,3,2)$	$x^3+y^4+z^4+w^6$	w	18	4
3	$(2,2,1,1)$	$x^2y+y^3+z^6+w^6$	z	10	8
4	$(4,4,3,1)$	$x^2y+y^3+z^4+w^{12}$	z	14	6
5	$(3,1,1,1)$	$x^2+y^6+z^6+w^6$	y	9	9
6	$(5,2,2,1)$	$x^2+y^5+z^5+w^{10}$	w	6	4
	$(5,2,2,1)$	$x^2+y^5+yz^4+w^{10}$	z	10	8
7	$(4,2,1,1)$	$x^2+y^4+z^8+w^8$	y	10	6
	$(4,2,1,1)$	$x^2+y^4+z^8+w^8$	w	10	8
8	$(6,3,2,1)$	$x^2+y^4+z^6+w^{12}$	y	12	6
	$(6,3,2,1)$	$x^2+y^4+z^6+w^{12}$	z	12	8
9	$(10,5,4,1)$	$x^2+y^4+z^5+w^{20}$	y	14	4
	$(10,5,4,1)$	$x^2+y^4+z^5+w^{20}$	w	14	4
10	$(6,4,1,1)$	$x^2+y^3+z^{12}+w^{12}$	z	10	8
12	$(9,6,2,1)$	$x^2+y^3+z^9+w^{18}$	w	6	2
13	$(12,8,3,1)$	$x^2+y^3+z^8+w^{24}$	z	14	6
16	$(8,7,6,3)$	$x^3+y^3w+z^4+w^8$	z	14	6
17	$(5,5,3,2)$	$x^2y+y^3+z^5+zw^6$	w	17	5
18	$(3,3,2,1)$	$x^2y+y^3+z^4w+w^9$	z	14	6
19	$(3,2,2,1)$	$x^2y+y^4+z^4+w^8$	z	10	8
23	$(5,3,2,2)$	$x^2z+y^4+z^6+w^6$	w	12	6
29	$(15,6,5,4)$	$x^2+y^5+z^6+yw^6$	w	18	4
31	$(12,5,4,3)$	$x^2+y^4z+z^6+w^8$	y	18	4
33	$(9,4,3,2)$	$x^2+y^4w+z^6+w^9$	y	14	6
	$(9,4,3,2)$	$x^2+y^4w+z^6+w^9$	z	10	6
36	$(10,5,3,2)$	$x^2+y^4+yz^5+w^{10}$	w	16	6
37	$(8,4,3,1)$	$x^2+y^4+yz^4+w^{16}$	z	10	8
39	$(9,5,3,1)$	$x^2+y^3z+z^6+w^{18}$	w	15	7
40	$(7,4,2,1)$	$x^2+y^3z+z^7+w^{14}$	w	7	3
41	$(12,7,3,2)$	$x^2+y^3z+z^8+w^{12}$	w	15	7
42	$(5,3,1,1)$	$x^2+y^3z+z^{10}+w^{10}$	w	11	9
44	$(8,5,2,1)$	$x^2+y^3w+z^8+w^{16}$	z	14	6
52	$(12,9,8,7)$	$x^3+y^4+xz^3+zw^4$	y	20	2
	$(12,9,8,7)$	$x^3+y^4+xz^3+zw^4$	w	19	3
75	$(11,5,4,2)$	$x^2+y^4w+z^5w+w^{11}$	y	13	5
84	$(9,7,6,5)$	$x^3+xz^3+y^3z+yw^4$	w	20	2

TABLE 10. $K3$ surfaces and their mirror partners

$S:Y\#$	$S:B\#$	weight	r	$S^\vee Y\#$	$S^\vee B\#$	$20-r$	weight for mirror
1		$(1,1,1,1)$	8	56		12	$(11,8,6,5)$
				73	27	12	$(25,10,8,7)$
4		$(4,4,3,1)$	10	4		10	$(4,4,3,1)$
5	1	$(3,1,1,1)$	1	52		19	$(12,9,8,7)$
6	2	$(5,2,2,1)$	6	26		14	$(9,5,4,2)$
				34	16	14	$(15,7,6,2)$
				76	12	14	$(13,6,5,2)$
8	41	$(6,3,2,1)$	3	64		17	$(10,7,4,5)$
9		$(10,5,4,1)$	10	9		10	$(10,5,4,1)$
				71		10	$(7,4,3,1)$
10	42	$(6,4,1,1)$	2	65		18	$(14,11,5,3)$
				46	29	18	$(53,22,6,5)$
				80	40	18	$(22,13,5,4)$
11	17	$(15,10,3,2)$	12	24		8	$(5,4,2,1)$
12	8	$(9,6,2,1)$	6	27		14	$(11,8,3,2)$
				49	25	14	$(21,14,5,2)$
13	45	$(12,7,3,1)$	8	20		12	$(9,8,6,1)$
				59		12	$(8,7,5,1)$
14	26	$(21,14,6,1)$	10	14	26	10	$(21,14,6,1)$
				28		10	$(10,7,3,1)$
				45	36	10	$(14,9,4,1)$
				51	47	10	$(18,12,5,1)$
20		$(9,8,6,1)$	12	17		8	$(12,8,3,1)$
				72		8	$(7,5,2,1)$
21		$(2,1,1,1)$	2	30	39	18	$(20,8,7,5)$
				86		18	$(9,7,5,4)$
22		$(6,5,3,1)$	10	22		10	$(6,5,3,1)$
24		$(5,4,2,1)$	8	11	18	12	$(15,10,3,2)$
25		$(4,3,1,1)$	8	43	46	12	$(18,11,4,3)$
				48	48	12	$(24,16,5,3)$
				88		12	$(11,9.5,2)$
26		$(9,5,4,2)$	14	6	2	6	$(5,2,2,1)$
27		$(11,8,3,2)$	14	12	8	6	$(9,6,2,1)$
28		$(10,7,3,1)$	10	14	26	10	$(21,14,6,1)$
				28		10	$(10,7,3,1)$
				45	36	10	$(14,9,4,1)$
				51	47	10	$(18,12,5,1)$
30	39	$(20,8,7,5)$	18	21		2	$(2,1,1,1)$
32	4	$(7,3,2,2)$	10	10	42	10	$(7,3,2,2)$
34	16	$(15,7,6,2)$	14	6	2	6	$(5,2,2,1)$
35	35	$(14,7,4,3)$	16	66		4	$(3,2,1,1)$
37	31	$(8,4,3,1)$	9	58		11	$(6,5,4,1)$
38	17	$(15,8,6,1)$	11	50	19	9	$(15,10,4,1)$
				82	11	9	$(11,7,3,1)$
39	7	$(9,5,3,1)$	9	60		11	$(7,6,4,1)$
40	5	$(7,4,2,1)$	7	81	14	13	$(13,8,3,2)$

Table 11. $K3$ surfaces and their mirror partners (continued)

$S:Y\#$	$S:B\#$	weight	r	$S^\vee:Y\#$	$S^\vee:B\#$	$20-r$	weight for mirror
42	3	(5,3,1,1)	3	68		17	(13,10,4,3)
				83	28	17	(27,18,5,4)
				92	23	17	(19,11,5,3)
43	46	(18,11,4,3)	16	25		4	(4,3,1,1)
45	36	(14,9,4,1)	10	14	26	10	(21,14,6,1)
				28		10	(10,7,3,1)
				45	36	10	(14,9,4,1)
				51	47	10	(18,12,5,1)
46	29	(33,22,6,5)	18	10	42	2	(6,4,1,1)
48	48	(24,16,5,3)	16	25		4	(4,3,1,1)
49	25	(21,14,5,2)	14	12	8	6	(9,6,2,1)
50	19	(15,10,4,1)	9	38	17	11	(15,8,6,1)
				77	13	11	(13,7,5,1)
51	47	(18,12,5,1)	10	14	26	10	(21,12,6,1)
				28		10	(10,7,3,1)
				45	36	10	(14,9,4,1)
				51	47	10	(18,12,5,1)
52		(12,9,8,7)	19	5		1	(3,1,1,1)
56		(11,8,6,5)	19	1		1	(1,1,1,1)
58		(6,5,4,1)	11	37	31	9	(8,4,3,1)
59		(8,7,5,1)	12	13	45	8	(12,8,3,1)
				72		8	(7,5,2,1)
60		(7,6,4,1)	11	39	7	9	(9,5,3.1)
64		(10,7,4,3)	17	7	30	3	(4,2,1,1)
65		(14,11,5,3)	17	10	42	3	(6,4,1,1)
66		(3,2,1,1)	4	35	35	16	(14,7,4,3)
68		(13,10,4,3)	17	42	3	3	(5,3,1,1)
71		(7,4,3,1)	10	9	34	10	(10,5,4,1)
				71		10	(7,4,3,1)
72		(7,5,2,1)	8	20		12	(9,8,6,1)
				59		12	(8,7,5,1)
73	27	(25,10,8,7)	19	1		1	(1,1,1,1)
76	12	(13,6,5,2)	14	6	2	6	(5,2,2,1)
77	13	(13,7,5,1)	11	50	19	9	(15,10,4,1)
				82	11	9	(11,7,3,1)
78	10	(11,6,4,1)	10	10	42	10	(11,6,4,1)
80	40	(22,13,5,4)	18	10	42	2	(6,4,1,1)
81	14	(13,8,3,2)	13	40	5	7	(7,4,2,1)
82	11	(11,7,3,1))	9	38	17	11	(15,8,6,1)
				77	13	11	(13,7,5,1)
83	28	(27,18,5,4)	17	42	3	3	(5,3,1,1)
86		(9,7,5,4)	18	21		2	(2,1,1,1)
87		(5,4,3,1)	10	87		10	(1,3,4,5)
92	23	(19,11,5,3)	17	42	3	3	(5,3,1,1)